



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : C07K 1/04, 7/02, 14/00, A61K 39/04, 39/02, 39/385	A1	(11) International Publication Number: WO 00/18791 (43) International Publication Date: 6 April 2000 (06.04.00)
(21) International Application Number: PCT/DK99/00510 (22) International Filing Date: 29 September 1999 (29.09.99) (30) Priority Data: PA 1998 01233 29 September 1998 (29.09.98) DK (71) Applicant (for all designated States except US): STATENS SERUM INSTITUT [DK/DK]; Artillerivej 5, DK-2300 København S (DK). (71)(72) Applicant and Inventor: HOLM, Arne [DK/DK]; Skodsborgparken 20, 2.tv., DK-2942 Skodsborg (DK). (72) Inventors; and (75) Inventors/Applicants (for US only): JØRGENSEN, Rikke, Malene [DK/DK]; Frem 31, DK-2800 Lyngby (DK). ØSTERGAARD, Søren [DK/DK]; Borrebyvej 21, DK-2700 Brønshøj (DK). THEISEN, Michael [DK/DK]; N.J. Fjords Allé 20, 4.th, DK-1957 Frederiksberg C (DK). (74) Agent: HOFMAN-BANG A/S; Hans Bekkevolds Allé 7, DK-2900 Hellerup (DK).		(81) Designated States: AE, AL, AM, AT, AT (Utility model), AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, CZ (Utility model), DE, DE (Utility model), DK, DK (Utility model), DM, EE, EE (Utility model), ES, FI, FI (Utility model), GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SK (Utility model), SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i>
(54) Title: LIGAND PRESENTING ASSEMBLY (LPA), METHOD OF PREPARATION AND USES THEREOF		
(57) Abstract The present invention relates to a method for preparing a Ligand Presenting Assembly (LPA), an LPA, an immunological composition and a vaccine. The invention further relates to a method for generating antibodies, a kit for use in diagnosis and use of an LPA for preparing a pharmaceutical composition.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav	TM	Turkmenistan
BF	Burkina Faso	GR	Greece		Republic of Macedonia	TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's	NZ	New Zealand		
CM	Cameroon		Republic of Korea	PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

LIGAND PRESENTING ASSEMBLY (LPA), METHOD OF PREPARATION AND USES THEREOF

The present invention relates to a method for preparing a
5 ligand presenting assembly (LPA), an LPA, an immuno-
logical composition and a vaccine. The invention further
relates to a method for generating antibodies, a kit for
use in diagnosis and use of an LPA for preparing a
pharmaceutical composition.

BACKGROUND OF THE INVENTION

Peptides having an amino acid sequence corresponding to a
part of the amino acid sequence of a protein or antigen
15 are often used in immunoassays for screening for anti-
bodies or preparation of vaccines.

The peptides or immunogenic compounds are often coupled
to various carriers in order to enhance the immunogenic
20 effect. One such system is multiple presentation of
antigenic peptides, which has long been recognised as
valuable means for amplifying peptide immunogens. Most
commonly an antigen is attached to a natural carrier such
as a protein, a carbohydrate, a lipid or a liposome to
25 provide a vaccine component or a synthetic sero-
diagnostic.

There are, however, a number of problems with these
products which may be related to the carriers. Since
30 these are isolated from natural sources they may not be
of uniform quality and may further contain harmful
contaminants being themselves antigenic. When used for
vaccines they may in particular result in fevers and
tissue swelling. In attempts to overcome certain of these
35 problems much research has in later years been focused on
antigen presentation using purely synthetic carriers.

Dendritic polymers are a such class of polymers or synthetic proteins. Such polymers have been described widely in the literature. The polymers are often referred
5 to as dendritic because of the similarity with a branched tree.

A multimerisation known as multiple antigen peptide (MAP) using a branched architecture with a controlled and
10 limited array of branches was described by Tam in PCT/-US90/02039 (ref. 1).

MAPs have applications as synthetic vaccines and as diagnostic tools as well as other biochemical uses
15 including immunoassays, serodiagnosis, epitope mapping, inhibitors, artificial proteins, intracellular delivery, affinity purification of antibodies and presentation of T-cell epitopes (ref. 2).

20 All branched peptides are similar in that they branch from a core matrix or template which provide them with different dendrimeric properties (ref. 3, ref. 4). Most commonly, lysine is used as amino acid in the core matrix because it has two ends, the α - and ϵ -amino groups,
25 available for branching reactions.

Different types of branching have been reviewed recently (ref. 5, ref. 2) including cascade, pennant or radial type of branching arrangement (Fig. 1A, B and C in ref.
30 5). A stem peptide carrier has been reported which is a short lipopeptide carrier, the stem peptide, which is useful for side-chain presentation of peptides (Fig. 2) (ref. 6).

35 The MAPs described above all have in common that an amino group is the starting point for the antigen

synthesis resulting in peptides which are attached to the core matrix by the C-terminal end and, thus, having a free N-terminal end.

5 The effectiveness of a B cell epitope on MAP to elicit humoral responses depends on the presence of a T cell recognition site located in the stretch of amino acids. While this is not always the case, chimeric B-T constructs have been made producing strongly enhanced
10 immune responses. Such MAPs are known as diepitope MAPs. The effectiveness of diepitope MAPs over monoepitope MAP has among other been studied in synthetic vaccine models of the human immunodeficiency virus (HIV) type 1. There are many possible ways of arranging B and T cell epitopes
15 on MAPs. Most commonly, they are linked in a tandem arrangement. High antibody titers were elicited by MAPs containing equimolar amounts of the T and B epitopes, while monoepitope MAPs and B cell epitope monomers showed no immunogenic effect in A/J mice used in the
20 experiments, cf. ref. 5. MAPs may be lipidated by standard peptide synthesis to produce a full range of response without any extraneous adjuvant (ref. 8, ref. 9).

25 As mentioned above, MAPs have a number of other applications. Important areas are immunoassays and serodiagnosis of e.g. malaria, cirrhosis, HIV-1, schistoma mansonii, systemic lupus erythematosus, and Epstein-Barr virus (EBV).

30

Most studies of MAPs have been carried out using tetra- or octameric MAP. However, it is generally found that with peptides composed of more than 15 amino acids, no real advantage is obtained using an octameric as compared
35 to a tetrameric MAP (ref. 2).

In general, two methods can be used for MAP synthesis, namely the direct route and the indirect route. In both cases, the C-terminal core is first assembled on a solid support using diprotected Boc-Lys(Boc) in the case of Boc chemistry, or Fmoc-Lys(Fmoc) in the case of Fmoc chemistry in order to obtain the desired degree of branching.

In the direct route, synthesis is continued by stepwise assembly of the particular peptide on the lysinyl core by the well-known Merrifield method. This stepwise method produces peptide antigens with a C to N orientation. B and T cell epitopes may be synthesised in tandem, or, alternatively, each epitope can be synthesised on the different lysine arms of the core using orthogonal deprotection methods. Such methods include i.a. Boc-Fmoc, and Fmoc-Aloc lysines. Although this approach for MAP synthesis is convenient, the quality of the final product may be questionable and problems of obtaining a well defined product have been discussed (ref. 10). Problems arise during chain assembly because MAPs are macromolecules where each single chain may interact with each other, preventing high efficiency couplings and leading to deletion sequences which require effort to purify. Furthermore, the characterisation of these products by, e.g. electrospray mass spectroscopy (ES-MS) is also difficult and as a result, multi-dendritic peptide immunogens have often been used without full characterisation. In the recent investigations by Keah et al. (ref. 10), a capping procedure after each amino acid coupling has been introduced in MAP synthesis, which improves the yield of the desired multi-dendritic immunogens. For analysis, an enzymatic (tryptic) digestion method with the MAP construct, followed by characterisation of the enzymatic digest by reversed

phase high-performance liquid chromatography-ES-MS method has been developed (ref. 10).

For many immunisations with MAP, the heterogeneity is not necessarily critical but for clinical applications, it is absolutely necessary to have chemically unambiguous MAPs without side products.

These considerations led to the development of indirect methods for MAP synthesis, in which methods purified unprotected peptide segments are coupled to the core matrix. Two general methods for this type of ligation have been developed which, respectively, are based on thiol and carbonyl chemistries.

The thiol chemistry may be carried out by incorporation of chloroacetyl group(s) on the lysine matrix and subsequent coupling to a purified, synthetic N-terminal cysteinyl peptide to yield a MAP with unambiguous structure (ref. 11, ref. 12). The reverse placement may be achieved with thiol on the core matrix by using an S-acetyl group on the lysinyl core and the haloacetyl group on the peptide (ref. 13). The use of disulphide bond formation for MAP synthesis is an alternative which is a well-known process in protein and peptide chemistry (ref. 2). A drawback in this type of chemistry is among others interference with cysteine present in the native antigen.

In the carbonyl chemistry, condensation takes place between a carbonyl group and a weak base. There are two types of weak bases which may be used for this reaction. The first includes hydroxylamines and hydrazines, and the second includes 1,2-disubstituted moieties such as 1,2-aminoethanthiol and 1,2-aminoethanol. The two last groups are found in N-terminal cysteine and threonine, respectively, and the condensation products are

thiazolidine and oxazolidine. The carbonyl group on the core matrix may be obtained by periodate oxidation of N-terminal serine, threonine or cysteine.

- 5 Several investigations along these approaches for MAP construction have been carried out to develop other methods for preparing peptide dendrimers providing flexibility for N or C attachment to the matrix core as reviewed by Tam (ref. 2). The processes may need
- 10 optimised conditions and a major concern by the use of thiols in the thiazolidine formation is a side reaction due to oxidation to disulphides, leading to impure products.
- 15 In the direct MAP synthesis route with stepwise assembly of the desired peptide on the lysinyl core a peptide antigen having C to N orientation is obtained. Such a MAP may be useful for ELISA applications, where antigen recognition by an antibody is directed at the N-terminal
- 20 or preferably close to the N-terminal, while recognition of the C-terminal may be hindered. This is e.g. the case with the dominant epitope of the Outer surface protein C (OspC) from *Borrelia burgdorferi sensu lato* causing Lyme borreliosis. Epitope mapping of OspC using sera from
- 25 patients with neuroborreliosis led to identification of one single major immunodominant epitope within the C-terminal ten amino acid residues, the decapeptide PVVAESPKKP. Peptide binding studies and alanine scanning revealed a critical role for the PKKP-sequence and in
- 30 particular its terminal carboxyl group for the binding of IgM antibodies from patients with Lyme borreliosis. Thus substitution of the C-terminal proline or replacement of the carboxy group with a carboxamido group greatly reduced the ability of the peptides to compete with
- 35 OspC19-207 for the binding of IgM antibodies in all five sera investigated. This significant fact clearly exclude

a MAP formed by stepwise synthesis on a lysine core as this results in C-terminal attachment to the core matrix.

5 From the above discussion, it is clear that a system providing a choice between C-, and C- and N-terminal presentation of sequences, and at the same time being easily performed and yielding improved products suitable for various applications, is needed.

10 SUMMARY OF THE INVENTION

The present invention provides novel methods for preparing LPAs enabling presentation of desired sequence(s). In a further aspect, the present invention
15 relates to a method for preparing LPAs enabling presentation of desired sequences(s) and chemical moieties.

In another aspect, the present invention relates to an
20 LPA obtainable by the method.

In yet a further aspect, the present invention relates to immunological compositions and vaccines, as well as a method of generating antibodies in an animal, including a
25 human being.

Furthermore, the present invention relates to the use of such LPAs in various applications and kits.

30 The present invention is described in detail below.

BRIEF DESCRIPTION OF THE FIGURES

Figure 1 shows the MALDI-TOF MS spectrum of the crude
35 product $\text{NH}_2\text{CH}(\text{CH}_2\text{CO-ProValValAlaGluSerProLysLysPro-OH})_2$

(also denoted $\text{NH}_2\text{CH}(\text{CH}_2\text{CO-Seq. ID 1-OH})_2$) [LPA-III] of Example 3.

Figure 2 shows the MALDI-TOF MS spectrum of the crude product $\text{NH}_2\text{CH}(\text{CH}_2\text{CO-ProValValAlaGluSerProLysLysPro-OH})_2$ (also denoted $\text{NH}_2\text{CH}(\text{CH}_2\text{CO-Seq. ID 1-OH})_2$) [LPA-III] of Example 3.

Figure 3 shows the HPLC chromatogram of crude H-ProVal-ValAlaGluSerProLysLysPro-OH, cf. Example 3.

Figure 4 shows the HPLC chromatogram of crude $\text{NH}_2\text{CH}(\text{CH}_2\text{CO-ProValValAlaGluSerProLysLysPro-OH})_2$ (also denoted $\text{NH}_2\text{CH}(\text{CH}_2\text{CO-Seq. ID 1-OH})_2$) [LPA-III] of Example 3.

Figure 5 shows the HPLC chromatogram of the HPLC purified product $\text{NH}_2\text{CH}(\text{CH}_2\text{CO-ProValValAlaGluSerProLysLysPro-OH})_2$ (also denoted $\text{NH}_2\text{CH}(\text{CH}_2\text{CO-Seq. ID 1-OH})_2$) [LPA-III] of Example 3.

Figure 6 shows the MALDI-TOF MS spectrum of the crude product H-Lys-NHCH($\text{CH}_2\text{CO-ProValValAlaGluSerProLysLysPro-OH})_2$ (also denoted H-Lys-NHCH($\text{CH}_2\text{CO-Seq. ID 1-OH})_2$) [LPA-IV] of Example 4.

Figure 7 shows the ELISA results from Example 13.

Figure 8 shows the ELISA results from Example 14.

Figure 9 shows the MALDI-TOF MS spectrum of the crude product $\text{CH}_2(\text{CH}_2\text{CO-}\beta\text{-Ala-}\beta\text{-AlaLysGluProAsnLysGlyValAsnPro-AspGluVal}\beta\text{Ala})_2$ also denoted $\text{CH}_2(\text{CH}_2\text{CO-}\beta\text{-Ala-}\beta\text{-Ala-Seq. ID 4-}\beta\text{-Ala})_2$ [LPA-VII] of Example 7.

Figure 10 shows the MALDI-TOF MS spectrum of the gel filtrated product $\text{HC}(\text{CH}_2\text{CO-LysGluProAsnLysGlyValAsnPro-})_2$

AspGluVal β Ala)₂COOH also denoted HC(CH₂CO-Seq. ID 4- β -Ala)₂COOH [LPA-VIII] of Example 8.

Figure 11 shows the HPLC chromatogram of the gel
5 filtrated product HC(CH₂CO-LysGluProAsnLysGlyValAsn-ProAspGluVal β Ala)₂COOH also denoted HC(CH₂CO-Seq. ID 4- β Ala)₂COOH [LPA-VIII] of Example 8.

Figure 12 shows the MALDI-TOF MS spectrum of the crude
10 product Fmoc-NHCH(CH₂CO-AspArgValTyrIleHisProPheHisLeu-NH₂)₂ also denoted Fmoc-NHCH(CH₂CO-Seq. ID 5-NH₂)₂ [LPA-IX] of Example 10.

Figure 13 shows the HPLC chromatogram of the crude
15 product Fmoc-NHCH(CH₂CO-AspArgValTyrIleHisProPheHisLeu-NH₂)₂ also denoted Fmoc-NHCH(CH₂CO-Seq. ID 5-NH₂)₂ [LPA-IX] of Example 10.

DETAILED DESCRIPTION OF THE INVENTION

20

In a first aspect, the present invention provides a novel method for preparing an LPA enabling presentation of desired sequence(s), which method comprises the steps of

25

(a) providing by solid phase synthesis or fragment coupling ligand(s) comprising the desired sequence(s), the ligand(s) being attached to a solid phase,

30

(b) if necessary, deprotecting any N-terminal amino groups while the ligand(s) are still attached to the solid phase,

35

(c) reacting the ligand(s) having unprotected N-terminal amino groups with an achiral di-, tri- or

tetracarboxylic acid so as to provide a construct having a ring structure, and

- 5 (d) cleaving the construct from the solid phase so as to provide an LPA comprising ligand(s) having free C-terminal groups.

10 In a further aspect, the present invention provides a method for preparing an LPA enabling presentation of desired sequence(s), or desired sequence(s) and chemical moieties, which method comprises the steps of

- 15 (c¹) if present, prior to step (d), deprotecting any N-protected amino groups originating from the carboxylic acid used in step (c),

- 20 (c²) continuing the solid phase synthesis or fragment coupling so as to provide ligand(s) comprising desired sequence(s) having at least one N-protected N-terminal amino group and/or attaching chemical moieties, and

- 25 (c³) deprotecting, if present, any N-terminal amino group(s) prior to or after step (d).

A unique feature of the method according to the present invention is the step providing a ring structure. Surprisingly, by the method according to the present invention it is possible to prepare very long ring systems interconnected by reaction with the achiral di-, tri- or tetracarboxylic acid. The ring structure formed between desired sequences further enables additional presentation of desired sequence(s) and chemical moieties. Thus, the method of the invention is easily performed and, as mentioned above, step (c¹) and (c²) enable further presentation of desired sequences and/or

30
35

chemical moieties. The desired sequences of step (c²) may be identical or different from the sequences of step (a). Thus, LPAs are suitable and very flexible systems for polyfunctional constructs, and furthermore, products of high purity are obtained.

Formation of double chain peptides having hemoregulatory activity is disclosed in WO 93/24523 (ref. 14). The two peptide chains are joined together at the C α -atoms of non-terminal amino acids in equivalent positions in each peptide chain via a carbon-carbon bond or via divalent bridging group -A- where -A- is a C₁₋₆ carbon chain. The double chain peptides are constructed from a single chain of the general formula R^a-R^b-R^c-R^d-(R^e)_n-R^f where R^a, R^b, R^c, and R^d represent selected amino acids and R^e is -NH-CH(R)-CO-, R^f represents lysine, arginine or glycine and n is 0 or 1. The bridging group A is attached to the C α -atoms of amino acids R^d and bridging is accomplished by acylating the synthesis resin bearing the desired N-deprotected C-terminal residue with one half equivalent of Fmoc-protected diamino dicarboxylic acid of such a type as 2,7-diaminosuberic acid or 2,9-diaminosebacic acid with the aid of coupling agents DCC (dicyclohexyl carbodiimide) and HOBt. After deprotection peptide synthesis is continued with the remaining amino acids depending on the sequence in question. It is a characteristic and a necessary condition of the employed amino acids in order to obtain the claimed biologically active compounds that they contain two chiral centres. Furthermore, the peptide chains which are assembled with the said diamino dicarboxylic acid are limited to two amino acids each according to the depicted formula.

In recent investigations coupling with half equivalent of protected diamino dicarboxyl acids with synthesis resin bearing an N-deprotected C-terminal residue with up to

three amino acids in the peptide chain has been reported (ref. 15, ref. 16). As to the conditions of the bridging technique it can be learned from the 1993 report by Alberts et al. (ref. 17) that bridging with half
5 equivalent 2,7-bis(Boc-amino) suberic acid coupled to one equivalent of lysine attached to the synthesis resin (2Cl-Z) is slow and takes place over up to 4 days. Thus, ring formation of longer amino acid sequences may be similarly or more difficult requiring activation
10 conditions which could prevent formation of well defined products with optically active bridging compounds.

The review by Tuchscherer and Mutter (ref. 18) deals with template-based de novo design of synthetic proteins (so-
15 called TASPs, template-assembled synthetic proteins). De novo design of polypeptide sequences with a three-dimensional structure necessary for many biological functions is limited by the complex folding process of the polypeptide sequences. The problem of folding can be
20 bypassed through constructing protein-like molecules with a built in device for intramolecular folding, namely proteins of non-natural chain architecture, i.e. TASPs. The concept is to use topological templates to introduce tertiary structures of the peptide sequences by directing
25 attached peptide blocks to a predetermined three-dimensional packing arrangement. This method provides synthetic proteins in which a predetermined backbone confirmation act as host for selective attachment of functional sites (e.g. amino acid side chains or
30 peptides) in spatially defined positions via a spacer group. An early example of this method is the use of β -D-glucose as a template (scaffold) for the design of a non-peptidic mimic of the somatostatin agonist SRIF. Covalent attachment of the amino acid side chains of Phe, Trp and
35 Lys to D-glucose derivatives resulted in a molecule recognised by G-protein coupled receptors. The concept

and purpose described in this reference is clearly different from the concept of the present invention. There is no mentioning of synthesis or assembly of ligands using achiral di-, tri- or tetracarboxylic acids.

5

In WO 94/02506 (ref. 19) MAP systems are disclosed. The MAPs are constructed with the following structure: Peptide antigen-core-linker-lipid anchor-resin. The core is based on lysine technology with four lysine arms onto which the peptide is assembled by conventional stepwise synthesis by reaction of a terminal functionality, e.g. a carboxylic acid group or an amine group. Different active substituents may be included in the molecule, thus providing MAPs with multiple antigenicities. In the document, the use of di-, tri- or tetracarboxylic acids is not mentioned or suggested. Thus, the present invention is clearly based on a different concept.

In WO 98/32469 (ref. 20) compounds comprising a linear, branched or dendrimeric polymer backbone are disclosed. At least one reporter molecule is linked thereto. The polymer backbone comprises a plurality of amine-containing acids. The compounds are useful as therapeutic and diagnostic agents, in particular in medical imaging techniques. The concept is based on the recognition that co-polymers of amino acids carrying or attached to one or more reporter molecules (e.g. chelating moieties or absorbers) are particularly suitable for diagnostic or therapeutic use. Dendrimeric polymers are the preferred backbone moieties. The compounds are formed from monomers which act as branching sites, and with each successive branching a new so-called generation is formed. The dendrimeric backbone molecule preferably comprises a multiplicity of native or non-native amino acid residues arranged so as to extend radially outwards from a central core moiety. The amino acids may be terminally bonded


directly or attached via a linking group to one or more reporter groups. Alternatively, these may be terminally branched by the addition of further amino acid residues. The compounds are prepared using conventional stepwise
5 solution synthesis. The use of di-, tri- or tetracarboxylic acids as in the present invention in the preparation is not disclosed or suggested, and, thus, the present invention is based on another principle. The structure of the resulting compounds are clearly
10 different from the LPAs of the present invention.

As mentioned previously, the publication by Tam (ref. 2) concerns design and immunological application of MAPs. The review focuses on the recent progress achieved in MAP
15 technology. I.a. cascade and pennant designs of core matrices for branched peptides are described including cascade and asymmetrical MAP with branching lysine, cascade and symmetrical MAP, pennant branched peptide, and polylysine as carrier. The review article does not
20 disclose synthesis using di-, tri- or tetracarboxylic acids.

The method of the present invention is based on assembly of two identical chains attached to a solid phase by
25 means of achiral dicarboxylic, tricarboxylic or tetracarboxylic acids. By using achiral compounds racemisation problems are avoided, thus allowing rigorous activation conditions for difficult reactions. With the exception of glycine all amino acids have a α -carbon atom
30 including the amino dicarboxylic acids aspartic and glutamic acid. The use of chiral acids leads to racemisation, i.e. in stead of obtaining a single product, a mixture of diastereomers will be obtained, the extent varying with reaction conditions. For clinical
35 applications, this is unacceptable and formation of clinically harmful side-products is a potential risk.

Racemisation problems arise upon activation of the carboxylic group of a chiral coupling agent, and in particular when the procedure is performed in the presence of a base, during long reaction times and/or
5 when the reaction is carried out at elevated temperatures.

Suitable achiral di-, tri- and tetracarboxylic acids to be used in the present method have the general formula
10



wherein n and m independently are an integer of from 1 to 20, X is HN, A and B independently are a substituted or
15 unsubstituted C₁₋₁₀ alkyl, a substituted or unsubstituted C₂₋₁₀ alkenyl, a substituted or unsubstituted cyclic moiety, a substituted or unsubstituted heterocyclic moiety, or a substituted or unsubstituted aromatic moiety, or A and B together form a substituted or un-
20 substituted cyclic moiety, a substituted or unsubstituted heterocyclic moiety, or a substituted or unsubstituted aromatic moiety, or

n and m are 0 or an integer of from 1 to 20, X is
25 H₂N(CR₂)_pCR, or RHN(CR₂)_pCR, wherein p is 0 or an integer of from 1 to 20, wherein each R is H, a substituted or unsubstituted C₁₋₁₀ alkyl, a substituted or unsubstituted C₂₋₁₀ alkenyl, a substituted or unsubstituted cyclic moiety, a substituted or unsubstituted heterocyclic
30 moiety, or a substituted or unsubstituted aromatic moiety, and A and B are both a substituted or unsubstituted C₁₋₁₀ alkyl, a substituted or unsubstituted C₂₋₁₀ alkenyl, a substituted or unsubstituted cyclic moiety, a substituted or unsubstituted heterocyclic
35 moiety, or A and B together form a substituted or unsubstituted cyclic moiety, a substituted or un-

substituted heterocyclic moiety, or a substituted or unsubstituted aromatic moiety, or

n and m are 0 or an integer of from 1 to 20, X is
5 HO(CR₂)_pCR, HS(CR₂)_pCR, halogen-(CR₂)_pCR, HOOC(CR₂)_pCR,
ROOC(CR₂)_pCR, HCO(CR₂)_pCR, RCO(CR₂)_pCR, or [HOOC(A)_n]-
[HOOC(B)_m]CR(CR₂)_pCR, wherein p is 0 or an integer of
from 1 to 20, each R independently is H, a substituted or
unsubstituted C₁₋₁₀ alkyl, a substituted or unsubstituted
10 C₂₋₁₀ alkenyl, a substituted or unsubstituted cyclic
moiety, a substituted or unsubstituted heterocyclic
moiety, or a substituted or unsubstituted aromatic
moiety, and A and B are both a substituted or un-
substituted C₁₋₁₀ alkyl, a substituted or unsubstituted
15 C₂₋₁₀ alkenyl, a substituted or unsubstituted cyclic
moiety, a substituted or unsubstituted heterocyclic
moiety, or A and B together form a substituted or un-
substituted cyclic moiety, a substituted or unsubstituted
heterocyclic moiety, or a substituted or unsubstituted
20 aromatic moiety, or

n and m are 0 or an integer of from 1 to 20, X is
H₂N(CR₂)_p, RHN(CR₂)_p, HO(CR₂)_p, HS(CR₂)_p, halogen-(CR₂)_p,
HOOC(CR₂)_p, ROOC(CR₂)_p, HCO(CR₂)_p, RCO(CR₂)_p or
25 [HOOC(A)_n][HOOC(B)_m], wherein p is 0 or an integer of
from 1 to 20, wherein each R independently is H, a
substituted or unsubstituted C₁₋₁₀ alkyl, a substituted
or unsubstituted C₂₋₁₀ alkenyl, a substituted or
unsubstituted cyclic moiety, a substituted or
unsubstituted heterocyclic moiety, or a substituted or
30 unsubstituted aromatic moiety, and A and B together form
a substituted or unsubstituted cyclic moiety, a
substituted or unsubstituted heterocyclic moiety, or a
substituted or unsubstituted aromatic moiety.

35

The term C₁₋₁₀ alkyl is intended to comprise straight or branched chain alkyl groups having 1-10 carbon atoms, e.g. methyl, ethyl, propyl, isopropyl, butyl, and tert-butyl.

5

The term C₂₋₁₀ alkenyl is intended to comprise straight or branched chain alkenyl groups having 2-10 carbon atoms, e.g. ethenyl, propenyl, isopropenyl, butenyl, and tert-butenyl.

10

The term cyclic moiety is intended to comprise e.g. cyclohexan, and cyclopentane.

15

The term aromatic moiety is intended to comprise e.g. phenyl.

20

The expression A and B forms a cyclic, heterocyclic or aromatic moiety is intended to comprise e.g. cyclohexan, piperidine, benzene, and pyridine.

25

By reaction with a carboxylic acid, a construct of the type

X(CO-sequence)₂-solid phase

is obtained, wherein X is as defined above.

30

By the term sequence, a peptide comprising naturally occurring and/or non-naturally occurring amino acids, a PNA-sequence, or a peptidomimetic is meant. Naturally occurring amino acids are L- and D-forms of the 20 amino acids found in nature. Non-naturally occurring amino acids are e.g. modified naturally occurring amino acids. The term sequence is further intended to comprise one or more of such sequences. Examples of suitable peptidomimetics are given by Marshall in ref. 21 and include

35

oligo(N-substituted glycines), oligo carbamates, oligo-sulfones, and oligosulfoxides. Peptidomimetics also include peptoides and peptomers.

5 The term chemical moieties is intended to comprise an entity enhancing the solubility or immunogenicity of the LPA, an entity for directing the LPA to its target or a marker.

10 The group X permits directly or indirectly continued stepwise synthesis or fragment coupling of the same sequence, or of one or more different sequences and/or chemical moieties. In the case of peptides, LPAs with N to C orientation, or with simultaneous N to C and C to N
15 presentation of the sequences are obtained. This is a unique feature of the method of the present invention. Continued synthesis or fragment coupling (fragment attachment) of a sequence different from the first such as a B or T cell epitope affords a chimeric product.

20

In the case where X comprises a temporarily protected amino function, synthesis or coupling can be carried out directly after deprotection. Suitable activation of all carboxy-containing groups providing effective formation
25 of the ring system can be ensured using half equivalent carboxylic acid. In case of tri- or tetracarboxylic acids, the activated carboxy group may further be derivatised with a diamine such as ethylenediamine or an amine suitably functionalised for further reactions such
30 as a mercapto-, an oxy-, an oxo or carboxyl group. In the case of a diamine, peptide synthesis or fragment coupling can be continued directly according to the desired sequence or chemical moiety. In a preferred embodiment, the Fmoc-protection strategy is used, but any amino
35 protection group may be used depending on the synthesis

or coupling strategy. Examples are the Boc-protection group strategy.

Since the continued stepwise synthesis or fragment
5 coupling is performed with one or in case of a
bifunctional chemical moiety such as lysine with two
amino groups, it has surprisingly been found that a much
better result can be obtained as compared to conventional
tetrameric or octameric MAP. This is believed to be due
10 to less interchain interaction. Furthermore, optimal
peptide synthesis procedures or coupling procedures can
be used for the single chains attached to the solid
phase, and their homogeneity can be verified prior to
forming the LPA. Cleavage from the solid phase and
15 simultaneous side-chain deprotection can be performed by
standard peptide synthesis procedures. A final LPA
product may thus be obtained having optimal and well
defined composition with possible by-products of less
than half the molecular weight of the target molecule.
20 Purification by standard chromatography methods such as
HPLC or gel filtration can easily be performed, if
desired or needed.

Favourable di-, tri- and tetracarboxylic acids for
25 providing the ring structure are e.g. imino diacetic
acid, 2-amino malonic acid, 3-amino glutaric acid, 3-
methylamino glutaric acid, 3-chloro glutaric acid, 3-
carboxymethyl glutaric acid, 3-methoxy-carbonyl glutaric
acid, 3-acetyl glutaric acid, glutaric acid,
30 tricarballic acid, 3,4-bis-carboxymethyl adipic acid,
4-(2-carboxyethyl)-pimelic acid, (3,5-bis-carboxymethyl-
phenyl)-acetic acid, 3,4-bis-carboxymethyl-adipic acid,
benzene-1,2,4,5-tetra carboxylic acid, 4-(3-carboxy-
allylamino)-but-2-enoic acid, 4,4-imino-dibenzoic acid,
35 1,4-dihydropyridine-3,5-dicarboxylic acid, 5-amino iso-

phthalic acid, 2-chloro malonic acid, 3-hydroxy glutaric acid, and benzene-1,3,5-tricarboxylic acid.

In the following, the individual steps of the method according to the present invention are described in further detail. Furthermore, apart from the general procedure, specific embodiments are given.

Step (a)

10

Solid phase synthesis has been known for a long time and can be carried out according to well-known standard procedures. Solid phase synthesis of peptides is e.g. described in WO 98/11125 (ref. 22). Solid phase synthesis of PNA sequences is e.g. described in WO 98/15648 (ref. 23).

Solid phase synthesis is generally carried out as described in the literature. However, a brief summary is given below.

The solid phase is preferably selected from functionalised resins such as functionalised polystyrene, polyacrylamide, polyethyleneglycol, cellulose, polyethylene, latex or dynabeads.

If desired, C-terminal amino acids may be attached to the solid support by means of a linker such as 2-Cl-trityl, 2,4-dimethoxy-4-hydroxybenzophenone, 4-(4-hydroxymethyl-3-methoxyphenoxy)-butyric acid (HMPB), 4-hydroxymethylbenzoic acid, 4-hydroxymethylphenoxyacetic acid (HMPA), 3-(4-hydroxymethylphenoxy)propionic acid or p-[(R,S)- α -[1-(9H-fluoren-9-yl)-methoxyformamido]-2,4-dimethoxybenzyl]-phenoxyacetic acid (AM).

35

The synthesis may be carried out batch-wise or continuously on an automated or semi-automated peptide synthesiser.

5 The individual coupling steps may be performed in the presence of a solvent, e.g. selected from acetonitrile, N,N-dimethylformamide (DMF), N-methylpyrrolidinone (NMP), dichloromethane (DCM), trifluoroethanol (TFE), ethanol, methanol, water, mixtures of the mentioned solvents with
10 or without additives such as perchlorate or ethylenecarbonate.

The individual couplings between two amino acids, an amino acid and the earlier formed sequence or fragment
15 may be carried out according to usual condensation methods such as the azide method, mixed acid anhydride method, symmetrical anhydride method, carbodiimide method, or active ester method using pentafluorophenyl (Pfp), 3,4-dihydro-4-oxobenzotriazin-3-yl (Dhbt), benzo-
20 triazol-1-yl (Bt), 7-azabenzotriazol-1-yl (At), 4-nitrophenyl, N-hydroxysuccinic acid imido esters (NHS), acid chlorides, acid fluorides, in situ activation by O-(7-azabenzotriazol-1-yl)-1,1,3,3-tetramethyluronium hexafluorophosphate (HATU), O-(benzotriazol-1-yl)-1,1,3,3-
25 tetramethyluronium hexafluorophosphate (HBTU), O-(benzotriazol-1-yl)-1,1,3,3-tetramethyluronium tetrafluoroborate (TBTU), benzotriazolyl-oxy-tris-(dimethylamino)-phosphonium hexafluorophosphate (BOP), benzotriazolyl-N-oxytripyrrolidinophosphonium hexafluorophosphate (PyBOP),
30 and 7-azabenzotriazolyl-N-oxytripyrrolidinophosphonium hexafluorophosphate (PyAOP).

Examples of temporary protection groups are Fmoc and Boc in the case of C α -amino protection. Examples of side-
35 chain protection groups are Boc, tBu, Z, and Cbz.

Examples of orthogonal protection groups are Fmoc, Boc, Mtt, Dde, All, ODmab, and Alloc.

5 Fmoc groups may be removed by means of an amine such as piperidine or diazabicyclo[5,4,0]undec-7-ene (DBU).

Side-chain protective groups may be removed by means of an acid such as TFA, TFMSA, HBr, HCl or HF.

10 To facilitate the reaction, the reaction mixture may be heated to a temperature of about 60°C.

Fragment coupling (fragment coupling or fragment condensation) may be performed according to standard
15 procedures, e.g. as described in ref. 24. Accordingly, fragments may be synthesised on a solid phase, cleaved from the solid phase with full preservation of protecting groups, purified and characterised. Synthesis of the fragments may be carried out as described above. Suitable
20 fragments may also be obtained by other techniques including enzymatic and recombinant techniques, and coupled as described above.

Step (b)

25

During solid phase synthesis or fragment coupling, N-terminal amino groups within the amino acid building blocks or side-chain amino groups of PNA moieties or peptidomimetics may be protected by suitable protecting
30 groups. Examples of temporary protection groups are Fmoc and Boc in the case of C α -amino protection. Examples of orthogonal protection groups are Mtt, Dde, All, ODmab, and Alloc.

35 If any protected amino groups are present, such may suitably be deprotected, while the sequences are still

attached on the solid phase. Suitable deprotecting agents are those usually used in the peptide chemistry and include piperidine, DBU or TFA in the case of temporary protection groups, and hydrazine, Pd/AcOH/NMM, or TFA in
5 the case of orthogonal protection groups.

Step (c)

A ring structure is provided using a di-, tri- or
10 tetracarboxylic acid. Formation of the ring structure is carried out under conditions enabling cyclisation of N-terminal amino groups of the desired sequences.

Suitable achiral di-, tri- and tetracarboxylic acids are
15 given above.

Suitably, the carboxylic acid may be used in an amount of approximately 0.4 to 0.6, preferably 0.5, equivalents carboxylic acid/peptide-resin.

20 Cyclisation may alternatively be carried out using 1:1 orthogonal N-terminal protection. After deprotection of 0.5 eq of the peptide-chain amino-groups the di-, tri- or tetracarboxylic acid in excess is activated and coupled.
25 After orthogonal deprotection cyclisation is achieved by activation of the second carboxylic acid group. Examples of orthogonal protection groups are Fmoc, Boc, Mtt, Dde, All, ODmab, and Aloc.

30 Activation may be carried out according to usual condensation methods such as the azide method, mixed acid anhydride method, symmetrical anhydride method, carbodi-imide method, or active ester method using Pfp, Dhbt, Bt, At, 4-nitrophenyl, NHS esters, acid chlorides, acid
35 fluorides, in situ activation by HATU, HBTU, TBTU, BOP, PyBOP, and PyAOP.

As already mentioned, the composition of the ligand may be verified and a sequence of the best possible purity may be achieved before the cyclisation procedure. If a
5 test of the so obtained product is found to contain monomeric substitution product, the procedure can be repeated to achieve a possible better yield. Alternatively, in the case where free peptide chains are present after cyclisation they may be terminated by e.g.
10 acetylation.

The reaction mixture may be heated up to about 60°C in order to facilitate the reaction. These harsh reaction conditions can be applied since an achiral carboxylic
15 acid is used. If a chiral carboxylic acid were used, elevated reaction temperatures would promote racemisation.

Step (d)

20 The resulting product, the construct, i.e. the LPA, is cleaved from the solid phase, whereby an LPA comprising ligands having free C-terminal groups are provided. Deprotection of any side-chain protecting groups can, if
25 desired, be carried out simultaneously with cleavage from the solid phase.

Cleavage from the solid phase can be carried out using well-known agents for this purpose such as TFA, TFMSA,
30 HBr, HCl, HF, ammonia, hydrazine, an alkoxide or a hydroxide. Alternatively, the sequence is cleaved from the solid phase by means of photolysis.

Furthermore, because the method of the invention provides
35 an LPA of approximately the double molecular weight,

standard chromatography methods such as HPLC or gel filtration can be used to obtain a single target LPA.

Step (c¹)

5

If a N-protected amino group originating from the carboxylic acid used in step (c) is present, such may be deprotected. Deprotection may be carried out using well-known conditions and agents. The amino group may suitably be protected by Fmoc or Boc. Suitable deprotecting agents are those usually used in the peptide chemistry and include piperidine, and DBU or TFA.

Step (c²)

15

Additional presentation of ligands comprising desired sequence(s) and/or chemical moieties may be provided by synthesis or coupling as described above.

Dicarboxylic acids

By the deprotection in step (c¹), a free amino group is obtained. The side chains of the sequence amino acid groups are preferably still fully protected.

Synthesis or fragment coupling may then be continued on the free amino group to provide a ligand comprising one or more desired sequences, said ligand having N-protected N-terminal amino group(s). Synthesis or coupling can be carried out using peptide chemistry, synthetic procedures appropriate for PNA, or procedures appropriate for peptidomimetics. Alternatively, synthesis or coupling may be carried out so as to incorporate a chemical moiety. Such chemical moieties may suitably be a fatty acid or another compound capable of improving the solubility of the LPA or immunogenicity (adjuvant effect), an entity suitable for directing the LPA to its target, a marker,

or any other group providing specific desirable properties. Lysine and amino hexanoic acid are examples of such other groups and may be used for chain extension.

5 Synthesis or fragment coupling may also be continued on the free amino group to provide a construct terminating with a diamino acid such as lysine, the α - and ϵ -amino groups of this diamino acid may be suitably protected (the protection groups being identical or different).
10 Suitable protection groups (orthogonal protection groups) are Fmoc, Boc, Mtt, Dde, All, ODmab, and Alloc. Subsequently, solid phase synthesis or fragment coupling may be carried out on each of the amino groups (i.e. both the α - and the ϵ -amino group), or only on one of the
15 amino groups (i.e. the α - or the ϵ -amino group) to provide a final bifunctional or trifunctional LPA. Solid phase synthesis may be carried out to provide sequences at both the α - and the ϵ -amino group, or only on the α - or the ϵ -amino group, and at the same time incorporating
20 a chemical moiety on the other. Alternatively, chemical moieties may be incorporated on both the α - and the ϵ -amino group.

Examples of markers as chemical moieties are
25 fluorophores, biotin, dinitro benzoic acid, haptens such as digoxigenin, enzymes such as HRP, alkaline phosphatases, and soya bean peroxidase, europium, and dyes such as rhodamines, cyanine dyes, coumarin, R-phycoerythrin (RPE) and allophycoerythrin.

30

Tri- and tetracarboxylic acids

Prior to step (d) described above, the free carboxylic group may be activated and functionalised by use of a suitable compound containing amino functions and/or thiol
35 and/or hydroxy functions such as 1,2-diamino ethane, 2-aminoethanthiol, and 2-aminoethanol.

In case of functionalising using a compound having a diamino function, synthesis or coupling may then be continued on the free amino group so as to provide a
5 desired sequence having a N-protected N-terminal amino group. Synthesis can be carried out using peptide chemistry or synthesis appropriate for PNA or procedures appropriate for peptidomimetics.

10 If the activated and functionalised construct terminates in an amino acid such as lysine, the α - and ϵ -amino groups may be suitably protected (the protection groups being identical or different). Suitable protection groups (orthogonal protection groups) are Fmoc, Boc, Mtt, Dde,
15 All, ODmab, and Alloc. Subsequently, synthesis or coupling may be carried out on each of the amino groups (i.e. both the α - and the ϵ -amino group), or only on one of the amino groups (i.e. the α - or the ϵ -amino group) so as to provide a final bifunctional or trifunctional LPA. Solid
20 phase synthesis or fragment coupling may be carried out to provide sequences at both the α - and the ϵ -amino group, or only on the α - or the ϵ -amino group and at the same time incorporating a chemical moiety on the other. Alternatively, chemical moieties may be incorporated at
25 both the α - and the ϵ -amino group. Examples of suitable chemical moieties are given above.

Step (c³)

30 Deprotection of any protected amino groups may be carried out prior to, subsequently to or simultaneously with the cleavage in step (d). Suitable cleavage agents are mentioned above under step (d).

35 As mentioned above, a remarkable and advantageous feature by the method provides an LPA with very well-defined

composition as optimised synthetic or coupling procedures can be used for preparation of the sequences.

5 The significant aspect of the use of amino or imino dicarboxylic acids in step (c) is the readily possibility of continued synthesis or coupling. After removal of the temporary protecting Fmoc-group in solid phase-attached and side-chain protected product, coupling with a desired activated and protected amino acid component may be
10 effected. A typical example is attachment of Fmoc-protected amino hexanoic acid followed by standard Fmoc-deprotection with piperidine and coupling with biotin with common activating agents. The resulting product is useful for e.g. ELISA applications with avidin or
15 streptavidin as coating agents.

The method provides i.a. LPAs presenting desired sequences with N to C orientation (using conventional terminology) (step (c)), and also simultaneously
20 sequences with C to N orientation (in the case of incorporation of ligands comprising amino acid sequences) (step (c²)).

In a preferred embodiment of the present invention, the
25 ligands comprising the desired sequence(s) comprise all or part of one or more B-cell epitopes, all or part of one or more T-cell epitopes, or all or part of one or more B- and T-cell epitopes (chimeras), or mimics thereof. The three-dimensional structure of a protein is
30 often such that amino acids, which are located distant from each other in the one-dimensional structure, are located near to each other in the assembled protein. Within the meaning of the present context, the expression epitope is intended to comprise the one- and three-
35 dimensional structure as well as mimics thereof. The term

is further intended to include discontinuous B-cell epitopes.

In accordance with the present invention, the desired sequences may be derived from viruses, bacteria, toxins, allergens, autoimmune system-related agents, cancer-related agents, cell adhesion molecules, neurotropic factors, fungi or parasites.

10 The peptide ligands may have substitutions, deletions or additions as long as the intended function of the ligand is sufficiently preserved. By the term substitution is meant substitution of one or more amino acids with another amino acids. By the term deletion is meant
15 removal of one or more amino acids from the sequence. By the term addition is meant insertion of one or more amino acids in the sequence. The sequence should have at least 40% identity, at least 50% identity, preferably at least 75% identity, and more preferably at least 95% identity,
20 with the sequence from which it is derived.

In accordance with the present invention, the desired sequence may preferably be derived from

25 viruses such as Myxoviruses e.g. influenza virus, Paramyxoviruses e.g. parainfluenza virus, mumps, measles, and Newcastle disease, Picornaviruses e.g. poliovirus, coxsackievirus, echovirus and rhinovirus, Reoviruses, Poxviruses e.g. as small pox virus, Vaccinia virus, and
30 cowpox virus, Papovaviruses e.g. polyoma virus, papilloma virus and SV-40, Adenoviruses, Astroviruses, SRSV (small round-structures virus), Epstein-Barr virus (EBV), Parvoviruses e.g. Human Parvovirus B19, Herpesviruses e.g. herpes simplex virus, cytomegalovirus (CMV),
35 Varicella virus (herpes zoster virus) and Pseudorabies virus, Syncytial viruses e.g. human syncytial virus

(HSV), Arboviruses e.g. yellow fever, and dengue fever, Leukoviruses e.g. Rous sarcoma virus, retroviruses e.g. HIV, and Hepatitis viruses e.g. Hepatitis A, Hepatitis B and Hepatitis C,

5

bacteria such as Mycobacterium spp. e.g. M. tuberculosis, M. bovis, M. africanum, M. microti, M. avium, M. intracellulare, M. kansasii, M. gordonae, M. paratuberculosis, and M. lepramurium, Borrelia spp. such as B. burgdorferi, in particular B. burgdorferi sensu stricto, B. garinii, B. afzelii, B. duttoni and B. recurrentis, Salmonella spp. e.g. S. typhimurium, and S. typhi, Treponema spp. e.g. T. pallidum, Leptospira spp., Campylobacter spp. e.g. C. jejuni, Helicobacter spp. e.g. H. pylori, Pseudomonas spp., Legionella spp., Neisseria spp. e.g. N. gonorrhoea and N. meningitidis, Chlamydia spp. e.g. C. trachomatis, C. pneumonia and C. psittae, Brucella spp., Enterobacter spp., Klebsiella spp., Yersinia spp., Vibrio spp. e.g. Vibrio cholerae, Gardnerella spp., Rickettsia spp., Clostridium spp. e.g. C. difficile and C. botulinum, Lactobacillus spp., Listeria spp., and Mycoplasma spp. e.g. M. pneumoniae and M. hominis,

25 toxins such as C. difficile toxin A and B, and C. tetani tetanospasmin,

allergens such as pollen allergens originating from the taxonomic order of Fagales, Oleales, Pinales, Poales, 30 Asterales, or Urticales, a mite allergen originating from Dermatophagoides, a cockroach allergen, and animal allergen originating from cat, dog or horse, a venom allergen originating from the taxonomic order of Hymenoptera, or a fungi allergens originating from the 35 genera Alternaria and Cladosporium,

auto-immune system-related compounds such as TNF α ,
cancer-related compounds,
5 cell adhesion molecules,
neurotropic factors,
fungi,
10 parasites such *Plasmodium* e.g. *P. falciparum*, *P. malariae*, *P. vivax*, and *P. ovale*, *Leishmania* e.g. *Leishmania donovani*, *L. brasiliensis*, *L. mexicana* and *L. peruviana*, or
15 a sequence homologous thereto.

Two bacterial species may be of particular interest, namely *Borrelia* spp. and *Mycobacterium* spp.

20 The tickborne spirochete *Borrelia burgdorferi* is the etiological agent of Lyme borreliosis, which is at present the most common vector-borne human disease in Europe and North America.

25 It is desirable to have access to an assay with a high diagnostic sensitivity already in the first stage of Lyme borreliosis, in order to diagnose and treat patients before they develop severe symptoms of the later stages
30 of Lyme borreliosis.

According to Western blot studies there are only two *B. burgdorferi* antigens that meet the essential criterion of eliciting an early and strong antibody response in the
35 majority of patients. These are the *B. burgdorferi* flagellum and the outer surface protein C (OspC). Whereas the performance of EIAs (enzyme immuno assays) using

purified native *B. burgdorferi* flagellum is well documented, the reported experience with OspC EIAs is still limited.

5 WO 97/42221 (ref. 25) discloses that the 4 terminal amino acids Pro-Lys-Lys-Pro are essential in the immune reactivity between sera from patients suffering from early borreliosis and various OspC derivatives. In order to obtain an effective diagnostic agent a polypeptide
10 fragment which contains carboxyterminally a peptide with the general formula $A^5-A^4-A^3-A^2-A^1$, where A^1 , and A^4 , independently from each other, designate a residue of an amino acid, wherein a nitrogen atom capable of forming part of a peptide bond is part of a ring structure; A^2
15 and A^3 , independently from each other, designate residues or a positively charged or polar amino acid; and A^5 designates residues of any amino acid, such that the peptide of the formula above has a degree of sequence identity of at least 60% with the amino acid residue
20 subsequence of Ser-Pro-Lys-Lys-Pro is preferred. Such polypeptide fragments may advantageously be used in preparing a multimeric structure according to the invention. Particular preferred OspC carboxyterminal peptides are Val-Ala-Glu-Ser-Pro-Lys-Lys-Pro, Val-Val-
25 Ala-Glu-Ser-Pro-Lys-Lys-Pro, and Pro-Val-Val-Ala-Glu-Ser-Pro-Lys-Lys-Pro.

In one embodiment of the present method, at least one of the sequences is derived from a sequence, wherein the C-
30 terminal amino acids are important for an immune response.

In another embodiment of the present method, the desired sequence is derived from the OspC protein of *Borrelia burgdorferi*, or is a homologous sequence capable of
35 reacting with anti-OspC antibodies, or is capable of provoking an immune response. In particular, the LPA

obtained provides a C-terminal presentation of the C-terminal sequence Pro-Lys-Lys-Pro of OspC.

5 In accordance with the invention, the presentation of a carboxyterminal OspC related peptide as described above can be combined with presentation of another sequence relevant for diagnosing Lyme borreliosis. In particular, for early diagnosis of Lyme borreliosis sequences comprising relevant epitopes from the flagellum are
10 relevant. Other relevant sequences might be derived from OspA, OspB, OspD, OspE, OspF, Erp proteins, BmpA (P39), P100, P35, P37 and DbpA. In particular, sequences derived from OspC, the flagellum, BmpA, (P39) or P100 are suitable sequences for diagnostic uses. Sequences derived
15 from OspA, OspC or DbpA are suitable vaccine candidates.

In yet another embodiment, the desired sequence is derived from the flagellum of *Borrelia burgdorferi* or is a sequence homologous thereto capable of reacting with
20 anti-flagellum antibodies

In yet another embodiment of the present invention, the LPA providing C-terminal presentation of sequences derived from OspC of *Borrelia burgdorferi* further
25 comprises desired sequence(s) derived from the flagellum of *Borrelia burgdorferi*.

In one embodiment of the present method, the desired sequence is derived from *Mycobacterium tuberculosis*. In a
30 further embodiment, the LPA of the invention provides C-terminal presentation of the desired sequences of OspC derived from *Borrelia burgdorferi*, and N-terminal presentation of desired sequence(s) derived from *Mycobacterium tuberculosis*. In particular, the desired
35 sequence derived from *M. tuberculosis* may comprise the sequence of amino acids number 51 to 70 of ESAT-6 (ESAT-6, 51-70 sequence protein) or the ESAT-6, 1-17 sequence

protein of *Mycobacterium tuberculosis* (cf. WO 95/01441 ref. 26).

The present invention also concerns an LPA obtainable by
5 the method defined above. The LPAs of the invention
enables presentation of desired sequence(s), or
presentation of desired sequence(s) and chemical
moieties. Such sequences are suitably selected from a
peptide comprising naturally occurring and/or non-
10 naturally occurring amino acids, a PNA sequence, or a
peptidomimetic. The chemical moieties are suitably an
entity enhancing the solubility or immunogenicity of the
LPA, or is an entity suitable for directing the LPA to
its target, or a marker. In particular, the chemical
15 moiety may be selected from fatty acids, antibodies or
peptides for directing the LPA to its target, biotin,
enzymes such as horse radish peroxidase, alkaline
phosphatase and soya bean peroxidase, or nucleic acid
sequences.

20

In one embodiment, the LPA of the invention comprises all
or part of one or more B cell epitopes, all or part of
one or more T cell epitopes, or all or part of one or
more B and T cell epitopes, or a mimic thereof. In
25 particular, at least one of the sequences may be derived
from a sequence, wherein the C-terminal amino acids are
important for an immunogenic response.

In another embodiment, the LPAs of the invention
30 comprises sequences which are derived from viruses,
bacteria, toxins, allergens, autoimmune system-related
compounds, cancer related compounds, fungi or parasites,
or a sequence homologous thereto.

35 In another embodiment, the LPAs of the invention are
such, wherein the desired sequence is derived from the

OspC protein of *Borrelia burgdorferi*. In particular, the LPA obtained may provide a C-terminal presentation of the C-terminal sequence Pro-Lys-Lys-Pro of OspC.

5 Examples of suitable LPAs are

[LPA-I]: FmocN(CH₂CO-ProValValAlaGluSerProLysLysPro-OH)₂
also denoted FmocN(CH₂CO-Seq. ID 1-OH)₂,

10 [LPA-II]: biotin-NH(CH₂)₅CON(CH₂CO-ProValValAlaGluSerPro-
LysLysPro-OH)₂ also denoted biotin-NH(CH₂)₅CON(CH₂CO-Seq.
ID 1-OH)₂,

[LPA-III]: NH₂CH(CH₂CO-ProValValAlaGluSerProLysLysPro-
15 OH)₂ also denoted NH₂CH(CH₂CO-Seq. ID 1-OH)₂, and

[LPA-IV]: H-Lys-NHCH(CH₂CO-ProValValAlaGluSerProLysLys-
Pro-OH)₂ also denoted H-Lys-NHCH(CH₂CO-Seq. ID 1-OH)₂.

20 In another embodiment of the LPA of the present
invention, the desired sequence is derived from
Mycobacterium tuberculosis. In a further embodiment of
the LPA, C-terminal presentation of the desired sequences
derived from OspC of *Borrelia burgdorferi*, and N-terminal
25 presentation of desired sequence(s) derived from
Mycobacterium tuberculosis are provided.

In particular, the LPA providing C-terminal presentation
of the C-terminal Pro-Lys-Lys-Pro of OspC of *Borrelia*
30 *burgdorferi*, and N-terminal presentation of ESAT-6
sequences from *M. tuberculosis* may be selected from

[LPA-V]: (HO-ProLysLysProSerGluAlaValValPro-COCH₂)₂CH-NH-
Lys(GlnLeuAlaAsnAsnLeuGluThrAlaThrAlaAspTrpLysGlnGln-
35 ValGlyGlnTyr-H)₂ also denoted (HO-Seq. ID 1-COCH₂)₂CH-NH-
Lys(Seq. ID 2-H)₂, and

[LPA-VI]: (HO-ProLysLysProSerGluAlaValValPro-COCH₂)₂N-Lys(AlaSerAlaAlaAlaGluIleGlyAlaPheAsnTrpGlnGlnGluThrMet-H)₂ also denoted (HO-Seq. ID 1-COCH₂)₂N-Lys(Seq. ID 3-H)₂.

Other examples of LPAs of the present invention are

[LPA-VII]: CH₂(CH₂CO-β-Ala-β-AlaLysGluProAsnLysGlyValAsn-ProAspGluValβAla)₂ also denoted CH₂(CH₂CO-β-Ala-β-Ala-Seq. ID 4-β-Ala)₂ of Chlamydia trachomatis DnaK 357-368,

[LPA-VIII]: HC(CH₂CO-LysGluProAsnLysGlyValAsnProAspGlu-ValβAla)₂COOH also denoted HC(CH₂CO-Seq. ID 4-βAla)₂COOH of Chlamydia trachomatis DnaK 357-368,

[LPA-IX]: Fmoc-NHCH(CH₂CO-AspArgValTyrIleHisProPheHisLeu-NH₂)₂ also denoted Fmoc-NHCH-(CH₂CO-Seq. ID 5-NH₂)₂ of Angiotensin-I,

[LPA-X]: Aloc-NHCH(CH₂CO-AspArgValTyrIleHisProPheHisLeu-NH₂)₂ also denoted Aloc-NHCH-(CH₂CO-Seq. ID 5-NH₂)₂ of Angiotensin-I, and

[LPA-XI]: Fmoc-AspProThrGlnAsnIleProProGly-NHCH(CH₂CO-AspArgValTyrIleHisProPheHisLeu-NH₂)₂ also denoted Fmoc-Seq. ID 6-NHCH(CH₂CO-Seq. ID 5-NH₂)₂ (Angiotensin I extended with Chlostridium thermosaccharolyticum peptide sequence 19-27).

In yet a further aspect, the present invention relates to an immunological composition for raising an immune response in an animal, including a human being comprising an LPA as defined above. In one embodiment, the immunological composition may comprise an LPA as defined above i.a. in combination with an adjuvant.

The immunological composition may suitably be in the form of a vaccine. Such a vaccine may suitably be for oral or parenteral, e.g. subcutaneous, intramuscular, intradermal, nasal, or pulmonary administration.

In another aspect, the present invention relates to a method for generating antibodies in an animal, including a human being, which method comprises administering to said mammal an antibody-generating amount of an LPA or an immunological composition as defined above.

Furthermore, the present invention relates to the use of an LPA as defined above for preparing a pharmaceutical composition for the treatment, alleviation or prophylaxis of diseases caused by viruses, bacteria, toxins, allergens, autoimmune system-related compounds, cancer related compounds, cell adhesion molecules, neurotropic factors, fungi or parasites.

Formulation of vaccines

Although it is preferred to insert a whole B- or T-cell epitope, it may in some cases be advantageous to insert a sequence comprising both the epitope as well as flanking regions from the protein from which the epitope is derived. The amino acids of a given epitope may be located near each other when the sequence is in the three-dimensional structure, but distant when the sequence is denatured. The sequence may be composed of the amino acids actually forming the epitope in the three-dimensional structure omitting all or part of the other amino acids.

According to the invention the LPAs may be used in vaccines against infections with pathogenic agents as

described above. Strategies in formulation development of vaccines based on the products obtained by the method of the present invention generally correspond to formulation strategies for any other protein-based drug product.

5 Potential problems and the guidance required to overcome these problems are dealt with in several textbooks. The use of an adjuvant, e.g., aluminium hydroxide, aluminium phosphate (Adju-Phos), calcium phosphate, muramyl dipeptide analogue, or some of the more recent

10 developments in vaccine adjuvants such as biodegradable microparticles and Iscoms is a formulation challenge familiar to a pharmaceutical scientist working in this area.

15 Preparation of the vaccines according to the invention which contain peptide sequences as active ingredients is generally well understood in the art, as exemplified by US 4 608 251 (ref. 27), US 4 601 903 (ref. 28), US 4 599 231 (ref. 29), US 4 599 230 (ref. 30), US 4 596 792 (ref.

20 31), and US 4 578 770 (ref. 32), all incorporated herein by reference. Typically, such vaccines are prepared as injectables either as liquid solutions or suspensions; solid forms suitable for solution in, or suspension in, liquid prior to injection may also be prepared. The

25 preparation may also be emulsified. The active immunogenic ingredient is often mixed with excipients which are pharmaceutically acceptable and compatible with the active ingredient. Suitable excipients are, for example, water, saline, dextrose, glycerol, ethanol, or the like,

30 and combinations thereof. In addition, if desired, the vaccine may contain minor amounts of auxiliary substances such as wetting or emulsifying agents, pH buffering agents, or adjuvants which enhance the effectiveness of the vaccines.

The vaccines are conventionally administered parenterally, by injection, for example, either subcutaneously or intramuscularly. Additional formulations which are suitable for other modes of administration include suppositories and, in some cases, oral formulations. For suppositories, traditional binders and carriers may include, for example, polyalkylene glycols or triglycerides; such suppositories may be formed from mixtures containing the active ingredient in the range of 0.5% to 10%, preferably 1-2%. Oral formulations include such normally employed excipients as, for example, pharmaceutical grades of mannitol, lactose, starch, magnesium stearate, sodium saccharine, cellulose, magnesium carbonate, and the like. These compositions take the form of solutions, suspensions, tablets, pills, capsules, sustained release formulations or powders and contain 10-95% of active ingredient, preferably 25-70%.

The LPA according to the invention may be formulated into the vaccine as neutral or salt forms. Pharmaceutically acceptable salts include acid addition salts (formed with the free amino groups of the peptide) and which are formed with inorganic acids such as, for example, hydrochloric or phosphoric acids, or such organic acids as acetic, oxalic, tartaric, mandelic, and the like. Salts formed with the free carboxyl groups may also be derived from inorganic bases such as, for example, sodium, potassium, ammonium, calcium, or ferric hydroxides, and such organic bases as isopropylamine, trimethylamine, 2-ethylamino ethanol, histidine, procaine, and the like.

The vaccines are administered in a manner compatible with the dosage formulation, and in such amount as will be therapeutically effective and immunogenic. The quantity to be administered depends on the subject to be treated, including, e.g., the capacity of the individual's immune

system to mount an immune response. Suitable dosage ranges are of the order of several hundred micrograms active ingredient per vaccination with a preferred range from about 0.1 μg to 1000 μg , such as in the range from about 1 μg to 300 μg , and especially in the range from about 10 μg to 50 μg . Suitable regimens for initial administration and booster shots are also variable but are typified by an initial administration followed by subsequent inoculations or other administrations.

The manner of application may be varied widely. Any of the conventional methods for administration of a vaccine are applicable. These are believed to include oral application on a solid physiologically acceptable base or in a physiologically acceptable dispersion, parenterally, by injection or the like. The dosage of the vaccine will depend on the route of administration and will vary according to the age of the person to be vaccinated and, to a lesser degree, the size of the person to be vaccinated.

Some of the LPAs according to the present invention may be sufficiently immunogenic in a vaccine, but for some of the others the immune response will be enhanced if the vaccine further comprises an adjuvant substance.

Various methods of achieving adjuvant effect for the vaccine include use of agents such as aluminium hydroxide or aluminium phosphate (alum, Adju-Phos), commonly used as 0.05 to 0.1 percent solution in phosphate buffered saline, admixture with synthetic polymers of sugars (Carbopol) used as 0.25 percent solution, aggregation of the protein in the vaccine by heat treatment with temperatures ranging between 70°C to 101°C for 30 second to 2 minute periods respectively. Aggregation by reactivating with pepsin treated (Fab) antibodies to

albumin, mixture with bacterial cells such as *C. parvum* or endotoxins or lipopolysaccharide components of gram-negative bacteria, emulsion in physiologically acceptable oil vehicles such as mannide mono-oleate (Aracel A) or
5 emulsion with 20 percent solution of a perfluorocarbon (Fluosol-DA) used as a block substitute may also be employed. According to the invention DDA (dimethyldi-octadecylammonium bromide) is an interesting candidate for an adjuvant, but also QuilA and RIBI are interesting
10 possibilities. Further possibilities are monophosphoryl lipid A (MPL), and muramyl dipeptide (MDP). Other suitable adjuvants are calcium phosphate, and muramyl dipeptide analogue. Some of the more recent developments in vaccine adjuvants such as biodegradable microparticles
15 and Iscoms may also suitably be used.

Another highly interesting possibility of achieving adjuvant effect is to employ the technique described in Gosselin *et al.*, 1992 (which is hereby incorporated by
20 reference). In brief, the presentation of a relevant antigen can be enhanced by conjugating such antigen to antibodies (or antigen binding antibody fragments) against the Fc γ receptors on monocytes/macrophages. Especially conjugates between antigen and anti-Fc γ RI have
25 been demonstrated to enhance immunogenicity for the purposes of vaccination.

Other possibilities involve the use of immune modulating substances such as lymphokines (e.g. IFN- γ , IL-2 and IL-
30 12) or synthetic IFN- γ inducers such as poly I:C in combination with the above-mentioned adjuvants.

In many instances, it will be necessary to have multiple administrations of the vaccine, usually not exceeding six
35 vaccinations, more usually not exceeding four vaccinations and preferably one or more, usually at least

about three vaccinations. The vaccinations will normally be at from two to twelve week intervals, more usually from three to five week intervals. Periodic boosters at intervals of 1-5 years, usually three years, will be
5 desirable to maintain the desired levels of protective immunity.

One reason for admixing the LPAs of the invention with an adjuvant is to effectively activate a cellular immune
10 response.

Diagnosis

When the LPAs are used for diagnosis, they may be applied
15 in various immunoassays. Samples to be tested may be obtained from e.g. body fluids such as blood, serum, cerebro spinal fluid (CSF), pleural, pericardial, synovial, pus and bone marrow, sputum, laryngeal swabs, gastric lavage, bronchial washings, biopsies, aspirates,
20 expectorates, urine, and tissue samples and further food samples, soil, and water samples as well as cultures thereof.

Apart from the use in vaccines, the LPA of the invention
25 may be used as a diagnostic tool. Other biochemical uses of the LPA include epitope mapping, inhibitors, artificial protein, intracellular delivery, and affinity purification of antibodies.

30 When the LPA is used as a diagnostic tool, a sample is brought into contact with the LPA under suitable conditions, and any reaction between the LPA and the target material is observed or measured. Suitable assays for diagnostic purposes include immunosorbent assays such
35 as ELISA, dot blot, flow cytometry and skin tests. Such

assay formats may be carried out by procedures well-known to the person skilled in the art.

5 The present invention further concerns a kit for use in the diagnosis of infections caused by viruses, bacteria, toxins, allergens, auto-immune system-related compounds, cancer related compounds, fungi or parasites, which kit comprises an LPA as defined above together with means for detecting or visualising binding between the LPA and the
10 substance to be detected. In one embodiment, the kit may be for use in diagnosing diseases caused by *Borrelia burgdorferi sensu lato*, and comprising an LPA as defined above together with means for detecting or visualising binding between the LPA and the substance to be detected.

15 The present invention is further illustrated by the following non-limiting examples.

EXAMPLES

20

General apparatus

HPLC conditions

HPLC was preformed on a Waters 600 E instrument equipped
25 with a Waters 996 Photodiode array detector with a Waters Radial Pak 8 x 100 mm C18 reversed-phase column. Buffer A was 0.1vol % TFA in water and buffer B 90 vol% acetonitrile, 9.9 vol% water and 0.1 vol% TFA. The Buffers were pumped through the column at a flow rate of
30 1.5 ml/min using a linear gradient from 0% - 70% B (20 min), linear gradient from 70 - 100% B (1 min), isocratic 100% B (5 min).

Mass spectroscopy

35 MALDI TOF spectra were obtained on a Fisons ToFSpec E instrument. As matrix were in all cases used α -cyano-4-

hydroxy-cinnamic acid. Molecular ions were recorded as M^+H^+ or M^+Na^+ and M^+K^+ adduct ions.

Amino acid analysis

- 5 Amino acid analysis was performed with a Waters PICOTAG system.

Amino acid sequencing

- 10 N-Terminal amino acid sequence determination was performed with Applied Biosystems 494 Protein Sequencer, Procise instrument.

Peptide synthesis

- 15 General procedures

Apparatus and synthetic strategy

- Peptides were synthesised batch-wise in a polyethylene vessel equipped with a polypropylene filter for filtration using 9-fluorenylmethyloxycarbonyl (Fmoc) or
20 tert-butyloxycarbonyl, (Boc) as N- α -amino protecting group and suitable common protection groups for side-chain functionalities.

Solvents

- 25 Solvent DMF (N,N-dimethylformamide, Riedel de-Häen, Germany) was purified by passing through a column packed with a strong cation exchange resin (Lewatit S 100 MB/H strong acid, Bayer AG Leverkusen, Germany) and analysed for free amines prior to use by addition of 3,4-dihydro-
30 3-hydroxy-4-oxo-1,2,3-benzotriazine (Dhbt-OH) giving rise to a yellow colour (Dhbt-O⁻ anion) if free amines are present. Solvent DCM (dichloromethane, analytical grade, Riedel de-Häen, Germany) was used directly without purification. Solvent NMP (N-methylpyrrolidone, Riedel
35 de-Häen) was used without purification.

Amino acids

Fmoc-protected amino acids and corresponding pentafluorophenyl (Pfp) esters were purchased from MilliGen (UK), NovaBiochem (Switzerland) and Bachem (Switzerland), and the Dhbt-esters from NovaBiochem (Switzerland) in suitable side-chain protected forms. Boc-protected amino acids were purchased from Bachem (Switzerland).

10 Coupling reagents

Coupling reagent diisopropylcarbodiimide (DIC) was purchased from Riedel de-Häen, Germany, and distilled prior to use, dicyclohexylcarbodiimide (DCC) was purchased from Merck-Schuchardt, München, Germany, and purified by distillation. O-benzotriazolyl-N,N,N',N'-tetramethyluronium tetrafluoroborate (TBTU) was purchased from PerSeptive Biosystems GmbH Hamburg, Germany, and benzotriazolyl-N-oxy-pyrrolidinium phosphonium hexafluorophosphat (PyBOP) and 7-azabenzotriazolyl-N-oxy-pyrrolidinium phosphonium hexafluorophosphat (PyAOP) from PerSeptive.

Linker

Linker 4-hydroxymethylphenoxyacetic acid (HMPA, Novabiochem, Switzerland) was coupled to the resin if required as a preformed 1-hydroxybenzotriazole (HObt) ester generated by means of DIC.

Solid phase

Peptides synthesised according to the Fmoc-strategy were synthesised on three different types of solid support using 0.05 M or higher concentrations of Fmoc-protected activated amino acid in DMF. 1) PEG-PS (polyethylene-glycol grafted on polystyrene; TentaGel S NH₂ resin, 0.27 mmol/g, TentaGel S RAM resin (s = 0.23 mmol/g), Rapp Polymere, Germany or NovaSyn TG resin, 0.29 mmol/g

(denoted s), Novabiochem, Switzerland); 2) PepSyn K (Kieselguhr supported polydimethylacrylamide resin functionalised with sarcosine methyl ester 0.11 mmol/g; MilliGen, UK); 3) Chlorotrityl resin, 0.3 - 0.9 mmol/g (Novabiochem). In the two first cases the first amino acid was coupled to the resin via linker HMPA except for TentaSel S RAM which is coupled directly affording peptide amides. In the third case the first amino acid was coupled directly to the chlorotrityl resin. Peptides synthesised according to the Boc-strategy were synthesised on a Merrifield-resin (polystyrene-divinylbenzene) with the first amino acid attached (Novabiochem).

15 Catalysts and other reagents

Diisopropylethylamine (DIEA), was purchased from Aldrich, Germany, ethylenediamine from Fluka, and piperidine from Riedel-de H  en, Frankfurt, Germany. 4-(N,N-dimethylamino)pyridine (DMAP) was purchased from Fluka, Switzerland and used as a catalyst in coupling reactions involving symmetrical anhydrides. 3,4-dihydro-3-hydroxy-4-oxo-1,2,3-benzotriazine (Dhbt-OH) was obtained from Fluka, Switzerland, 1-hydroxybenzotriazole (HObt) from NovaBiochem, Switzerland and 1-hydroxy-7-azabenzotriazole (HOAt) from PerSeptive. Biotin was purchased from Sigma. Scavengers Triisopropylsilan, ethanedithiol and thioanisol was purchased from Aldrich, Germany.

General coupling procedures

30 The first amino acid was coupled as a symmetrical anhydride in DMF generated from the appropriate N-  -protected amino acid and DIC. The following amino acids were coupled as Pfp- or Dhbt-esters or as preformed HOBT or HOAt esters made from appropriate N-  -protected amino acids and HObt or HOAt by means of DIC or TBTU in DMF
35 unless otherwise stated.

Deprotection of the N- α -amino protecting group

Deprotection of the Fmoc group was performed by treatment with 20% piperidine in DMF (1x3 and 1x7 min) followed by wash with DMF until no yellow colour (Dhbt-O-) could be detected after addition of Dhbt-OH to the drained DMF. Deprotection of the Boc group was performed by treatment with 50% TFA in DCM (v/v) 1x1.5 min and 1x20 min followed by washing 6x9 min each with DCM, neutralisation with 10% triethylamine in DCM (v/v) 2x1.5 min each, followed by 6x9 min washing with DCM.

Cleavage from solid support

Cleavage of peptide and N-terminal product from solid phase and simultaneous side-chain deprotection were done with acid. Peptides were cleaved from the resins by treatment with 95% trifluoroacetic acid (TFA, Halocarbon Products Corporation, USA; Biesterfeld & Co. Hamburg, Germany)-water v/v or with TFA (trifluoroacetic acid)-H₂O-ethanedithiole-thioanisole (90:5:3:2 vol./vol./vol./-vol.) or with TFA-TIS and other scavengers at room temperature for 2 h. The filtered resins were washed with 95% TFA-water and filtrates and washings concentrated under reduced pressure. The residue was treated with ether and the precipitate freeze dried from acetic acid-water. The crude freeze dried product was analysed by HPLC and identified by MALDI-TOF MS.

Preformed symmetrical anhydride (PSA)

6 eq. N- α -amino protected amino acid was dissolved in DCM and cooled to 0°C. DCC (3 eq.) was added and the reaction continued for 10 min to yield PSA. The solvent was removed in vacuo and the remanens dissolved in DMF. The solution was filtered and immediately added to the resin followed by 0.1 eq. of DMAP.

Estimation of the coupling yield of the first N- α -amino protected amino acid

3-5 mg dry Fmoc-protected peptide-resin was treated with 5 ml 20% piperidine in DMF for 10 min at room temperature and the UV absorption for the dibenzofulvene-piperidine adduct was estimated at 301 nm. The yield was determined using a calculated extension coefficient e_{301} based on a Fmoc-Ala-OH standard.

10 In case of Boc-protection, the coupling was estimated according to the ninhydrin-method after removal of the Boc-group (Sarin et al., 1981, ref. 7).

Peptide synthesis on PepSyn K resin

15 Dry PepSyn K (ca 500 mg), was covered by ethylenediamine and left at room temperature over night. The resin was drained and washed with DMF 10 x 15 ml, 5 min each. After draining, the resin was washed with 10% DIEA in DMF v/v (2x15 ml, 5 min each) and finally washed with DMF until
20 no yellow colour could be detected by addition of Dhbt-OH to the drained DMF. 3 eq. HMPA 3 eq. HOBT and 3 eq. DIC was dissolved in 10 ml DMF and left for activation for 10 min, after which the mixture was added to the resin and the coupling continued for 24 h. The resin was drained
25 and washed with DMF (10x15 ml, 5 min each), and the acylation was checked by the ninhydrin test. The first amino acid was coupled as the preformed symmetrical anhydride (see above), and the coupling yields estimated as described above. It was in all cases better than 70%.
30 The remaining amino acids according to the sequence were coupled as described above. The couplings were continued for 2 h unless otherwise specified. Excess reagent was then removed by DMF washing. After completed synthesis the peptide-resin was washed with DMF and diethyl ether
35 and dried in vacuo.

Peptide synthesis on PEG-PS

TentaGel S NH₂ or NovaSyn TG resin (250 mg, 0.27-0.29 mmol/g) was placed in a polyethylene vessel equipped with a polypropylene filter for filtration. The resin was
5 swelled in DMF (5ml), and treated with 20% piperidine in DMF to secure the presence of non-protonated amino groups on the resin. The resin was drained and washed with DMF until no yellow colour could be detected after addition of Dhbt-OH to the drained DMF. HMPA (3 eq.) was coupled
10 as a preformed HObt-ester as described above and the coupling was continued for 24 h. The resin was drained and washed with DMF (5x5 ml, 5 min each) and the acylation checked by the ninhydrin test. The first amino acid was coupled as a preformed symmetrical anhydride as
15 described above. The coupling yields of the first Fmoc-protected amino acids were estimated as described above. It was in all cases better than 70%. The following amino acids according to the sequence were coupled as preformed Fmoc-protected HObt esters (3 eq.) as described above.
20 The couplings were continued for 2 h, unless otherwise specified. The resin was drained and washed with DMF (5x5 ml, 5 min each) in order to remove excess reagent. All acylations were checked by the ninhydrin test performed at 80 °C. After completed synthesis the peptide-resin was
25 washed with DMF (3x5 ml, 5 min each), DCM (3x5 ml, 1 min each) and finally diethyl ether (3x5 ml, 1 min each) and dried in vacuo.

Peptide synthesis on chlorotrityl resin

30 The chlorotrityl resin (approximately 0.8 mmol/g, Nova-Biochem) and the first Fmoc-protected amino acid in the sequence was dried in vacuo overnight before use. Solvents, DCM and DMF, was dried over molecular sieves (4 Å, 24h) in a round bottomed flask for derivatisation
35 dried at 100°C. The resin was placed in the flask under argon and the Fmoc-amino acid (10 eq compared to the eq

of resin) was dissolved in DCM (ca. 0.5 M) with a small amount of DMF added if necessary. The solution was added to the resin followed by addition of DIEA (20 eq) and the mixture left for 30 min with stirring. The resin was
5 transferred to a syringe fitted with a filter and washed with DCM/MeOH/DIEA (17:2:1) 3 times to cap remaining Cl, followed by washing with 3xDCM and 3xDMF. A small amount of resin, washed with DCM and MeOH prior to drying in a dessicator, was used for determination of substitution as
10 described above by formation of dibenzofulvene-piperidine. However, treatment with piperidine was extended to at least 3-4 h prior to analysis because of slow deprotection of the Fmoc-group. After deprotection of the Fmoc-group using a prolonged deprotection, the
15 remaining amino acids according to the sequence were coupled as described above. The couplings were continued for 2 h unless otherwise specified. Excess reagent was then removed by DMF washing. After completed synthesis the peptide-resin was washed with DMF and diethyl ether
20 and dried in vacuo.

Fmoc-iminodiacetic acid (Fmoc-IDA)

The product was prepared by the following method: iminodiacetic acid (2.5 g, 0.0188 mol, Aldrich) was dissolved
25 in 10% aqueous sodium carbonate, pH 9.5 (150 ml). Fmoc-Cl (4.85 g, 0.0188 mol, Bachem) was dissolved in tetrahydrofuran (50 ml) and added dropwise over 30 min to the iminodiacetic acid solution. The mixture was left for 5 hr at room temperature, washed twice with ethyl acetate,
30 acidified with aqueous HCl to pH 2 and Fmoc-iminodiacetic acid was extracted twice with ethylacetate. The combined extracts were washed twice with aqueous HCl (pH 2) followed by washing with brine and then concentrated in vacuo. Pentane was added to turbidity and the
35 crystallisation was allowed to proceed for 10 min on an ice bath. The Fmoc-iminodiacetic acid was collected by

filtration and the filtrate washed with cold ethylacetate/pentane and dried to give 5.61 g (84% yield). According to HPLC the purity was >99%. Mp. 212-214 °C. Literature. 213 - 215.5 °C.

5

**N-(Fluoren-9-ylmethoxycarbonyl)-2-aminomalononic acid
(Fmoc-2-AMA)**

Diethyl 2-aminomalonate (Sigma, 200 mg, 0.95 mmol) was dissolved in an aqueous solution of sodium carbonate (10%, 6 ml) and the mixture refluxed and stirred for 3h after which TLC (EtOH:AcOH (4:1) and development with ninhydrin) showed complete hydrolysis of the diethyl ester. To the reaction mixture was added 1.2 eq of O-(Fluoren-9-ylmethoxycarbonyl)-hydroxysuccinimide (Nova-
15 Biochem, Fmoc-OSu, 380 mg, 1.13 mmol) dissolved in dioxane (2 ml) over a period of 15 min and the solution stirred at room temperature overnight. Upon dilution with dioxane, the reaction mixture was acidified with HCl (1 M) to pH 5 and extracted with dichloromethane. The
20 aqueous phase was further acidified with HCl (1 M) to pH 2 upon which the product precipitated. The aqueous phase was added ethyl acetate and stirred for 10 min. prior to extraction with ethyl acetate. The organic phase was washed with water, dried over MgSO₄ and concentrated in
25 vacuo. The product, was crystallised from ethyl acetate - heptane with an overall yield of 135 mg (42%), mp. approximately 164°C, decomp. FAB-MS: Found 342; calculated 341.3. RP-HPLC analysis, t_R 19 min, using above described conditions, revealed a minimal content of the
30 decarboxylated product, Fmoc-Gly-OH(< 3%).

**N-(Fluoren-9-ylmethoxycarbonyl)-3-aminoglutaric acid
(Fmoc-3-AGA)**

3-Aminoglutaric acid (Sigma, β-glutamic acid, 30 mg, 0.20
35 mmol) was dissolved in 10% sodium carbonate/dioxane (1 ml/0.5 ml) and added Fmoc-OSu (83 mg, 0.25 mmol)

dissolved in dioxane (1 ml) over a period of 15 min. After stirring at room temperature overnight, the reaction mixture was diluted with water and concentrated in vacuo to remove dioxane prior to extraction with DCM.

5 The aqueous phase was separated, acidified with HCl (1 M) to pH 2-3 and the precipitated product extracted into ethyl acetate. The organic phase was washed with water, dried over MgSO₄ and concentrated in vacuo. The product was crystallised from dioxane-water yielding 47 mg (63%),

10 mp. 189-190 °C, RP-HPLC analysis, t_R 21 min using above described conditions. FAB-MS: Found 370; calculated 369.4

Allyloxycarbonyl-3-aminoglutaric acid (Aloc-3-AGA)

3-Aminoglutaric acid (107 mg, 0.73 mmol) was dissolved in

15 H₂O-dioxan (2.5 ml/2 ml). Triethylamine (0.2 ml, 2 eq.) and diallyl pyrocarbonate (0.12 ml, 1 eq., Lancaster, Mühlheim am Main, Germany) was added and the mixture heated under reflux over night. After evaporation of dioxan *in vacuo* the mixture was acidified with citric

20 acid (10%, 3 ml) and extracted with ethyl acetate (3x5 ml). The organic phase was washed with water, dried over MgSO₄ and finally concentrated *in vacuo*. Trituration with pentane yielded 76 mg (45%). M.p. 117-118°C. FAB-MS: Calc. for C₉H₁₃NO₆, 231.2 found 232.0.

25

1-(4,4-Dimethyl-2,6-dioxo-cyclohex-1-ylidene)ethyl-β-Ala (Dde-β-Ala)

β-Ala (100 mg, 1.12 mmol) was dissolved in EtOH (98%, 8 ml). 2-Acetyldimedone (614 mg, 3.36 mmol, NovaBiochem)

30 and triethylamine (0.24 ml, 3.36 mmol) was added and the mixture was refluxed for 6 hours. After continued stirring over night, the mixture was concentrated to give an oil. Dissolution in ethyl acetate (5 ml) was followed by neutralisation with citric acid (10%, 3x5 ml). The

35 organic phase was washed with water, dried over MgSO₄ and finally concentrated *in vacuo*. Re-crystallisation from

the ethyl acetate gave 180 mg (63%). M.p. 213-214°C. FAB-MS: Calc. for $C_{13}H_{19}NO_4$, 253.3 found 254.0.

EXAMPLE 1

5

Synthesis of FmocN(CH₂CO-ProValValAlaGluSerProLysLys-Pro)₂-chlorotrityl resin and FmocN(CH₂CO-ProValValAlaGluSerProLysLysPro-OH)₂ (IDA-synthesis of Seq. ID 1)

10 Product: FmocN(CH₂CO-ProValValAlaGluSerProLysLysPro-OH)₂ also denoted FmocN(CH₂CO-Seq. ID 1-OH)₂ [LPA-I]

The antigenic C-terminal sequence of OspC from *Borrelia burgdorferi* ProValValAlaGluSerProLysLysPro (Seq. ID 1) was assembled on chlorotrityl resin (s = 0.8 mmol/g) as described above and tested for homogeneity. The peptide moiety ProValValAlaGluSerProLysLysPro has a molecular weight of 1035 Dalton and a corrected substitution of 0.44 is used for samples of the peptide-resin. 100 mg of the peptide-resin was suspended in DMF and deprotected with 20% piperidine-DMF. 0.55 eq Fmoc-iminodiacetic acid (i.e. 10% excess; $0.1 \times 1.1 \times 0.05 \times 0.44 \times 355 = 8.6$ mg; Mw 355), 3.3 eq TBTU ($0.1 \times 1.1 \times 3 \times 0.44 \times 321 = 46.6$ mg; i.e. 3 times excess compared to the amount carboxy groups on Fmoc-iminodiacetic acid), 3.3 eq HOBT ($0.1 \times 1.1 \times 3 \times 0.44 \times 135 = 19.6$) and 4.95 eq DIEA ($0.1 \times 1.1 \times 4.5 \times 0.44 \times 174 = 38$ ml; standard amount of DIEA is 1.5×TBTU-eq.) was added to the resin and left overnight. Excess reagents were removed, the resin was washed with DMF, followed by ether and dried. The resulting dimeric compound, FmocN(CH₂CO-ProValValAlaGluSerProLysLysPro-OH)₂, was cleaved from the resin with 95% aqueous TFA. MALDI TOF MS: Calculated for $C_{115}H_{177}N_{25}O_{32}$, 2421.8; found 2423.2. The product contains only a small amount of the side-product FmocN(CH₂CO-ProValValAlaGluSerProLysLysPro-OH)CH₂COOH

15
20
25
30
35

(Calculated for $C_{67}H_{97}N_{13}O_{19}$, 1388.6; found 1389.0) which is essentially absent upon repeated coupling.

EXAMPLE 2

5

Synthesis of biotin-NH(CH₂)₅CON(CH₂CO-ProValValAlaGluSerProLysLysPro-OH)₂ (IDA derivative of Seq. ID 1 extended with 6-amino-hexanoic acid and biotinylated)

10 Product: biotin-NH(CH₂)₅CON(CH₂CO-ProValValAlaGluSerProLysLysPro-OH)₂ also denoted Biotin-NH(CH₂)₅CON(CH₂CO-Seq. ID 1-OH)₂ [LPA-II]

FmocN(CH₂CO-ProValValAlaGluSerProLysLysPro)₂-chlorotrityl
15 resin was deprotected with 20% piperidine-DMF and coupled with Fmoc-6-amino-hexanoic acid activated with TBTU. The coupling was followed by exhaustive acetylation with acetic anhydride. After deprotection with 20% piperidine-DMF, biotin (C₁₀H₁₆N₂O₃S) was coupled, activated with
20 TBTU. The product was cleaved from the resin with 95% aqueous TFA. MALDI TOF MS analysis showed the target compound as the major product (calculated for C₁₁₆H₁₉₂N₂₈O₃₃S, 2539.0; found 2536.3) containing some side-product, biotin-NH(CH₂)₅CON(CH₂CO-ProValValAlaGluSerProLysLysPro-OH)CH₂COOH (calculated for C₆₈H₁₁₂N₁₆-O₂₀S, 1505.8; found 1504.1).
25

EXAMPLE 3

30 Synthesis of FmocNHCH(CH₂CO-ProValValAlaGluSerProLysLysPro)₂-chlorotrityl resin and NH₂CH(CH₂CO-ProValValAlaGluSerProLysLysPro-OH)₂ (AGA-3-derivative of Seq. ID 1)

Product: NH₂CH(CH₂CO-ProValValAlaGluSerProLysLysPro-OH)₂
35 also denoted NH₂CH(CH₂CO-Seq. ID 1-OH)₂ [LPA-III]

The resin derivative was assembled in the same way as described above for the iminodiacetic acid product using N-(Fluoren-9-ylmethoxycarbonyl)-3-aminoglutaric acid. The resulting Fmoc-NHCH(CH₂CO-ProValValAlaGluSerProLysLys-Pro)₂-chlorotrityl resin was deprotected with 20% piperidine-DMF and NH₂CH(CH₂CO-ProValValAlaGluSerProLysLysPro-OH)₂ was obtained by cleavage from the resin with 95% aqueous TFA. MALDI-TOF MS: Calculated for C₁₀₁H₁₆₉N₂₅O₃₀, 2213.6; found 2215.2 (Figures 1 and 2). The higher molecular ions are the sodium (2238.0) and potassium ion adducts (2254.3) respectively. The product contains only a very small amount of lower molecular weight products as evidenced from the spectra. HPLC of crude H-ProValValAlaGluSerProLysLysPro-OH and crude NH₂CH(CH₂CO-ProValValAlaGluSerProLysLysPro-OH)₂ is shown in Figures 3 and 4, respectively. The HPLC of the HPLC-purified LPA-III is shown in Figure 5.

EXAMPLE 4

Synthesis of Fmoc-Lys(Fmoc)-NHCH(CH₂CO-ProValValAlaGluSerProLysLysPro)₂-chlorotrityl resin and H-Lys-NHCH(CH₂CO-ProValValAlaGluSerProLysLysPro-OH)₂ (AGA-3-derivative of Seq. ID 1 extended with di-Fmoc-lysine)

Product: H-Lys-NHCH(CH₂CO-ProValValAlaGluSerProLysLysPro-OH)₂ also denoted H-Lys-NHCH(CH₂CO-Seq. ID 1-OH)₂ [LPA-IV]

The above synthesised NH₂CH(CH₂CO-ProValValAlaGluSerProLysLysPro)₂-chlorotrityl resin was coupled with di-Fmoc-lysine in 3 times excess with TBTU as coupling agent. After Fmoc-deprotection with 20% piperidine-DMF, H-Lys-NHCH(CH₂CO-ProValValAlaGluSerProLysLysPro-OH)₂ was cleaved from the resin with 95% aqueous TFA and precipitated

with ether. MALDI TOF MS analysis: Calculated for $C_{107}H_{181}N_{27}O_{31}$, 2341.8; found 2343.2 (Figure 6).

EXAMPLE 5

5

Synthesis of (HO-ProLysLysProSerGluAlaValValPro-COCH₂)₂-CH-NH-Lys(GlnLeuAlaAsnAsnLeuGluThrAlaThrAlaAspTrpLysGln-GlnValGlyGlnTyr-H)₂(AGA-3-derivative of Seq. ID 1 extended with bis-ESAT-6, 51-70 sequence (Seq. ID 2), on lysine)

10

Product (HO-ProLysLysProSerGluAlaValValPro-COCH₂)₂CH-NH-Lys(GlnLeuAlaAsnAsnLeuGluThrAlaThrAlaAspTrpLysGlnGlnVal-GlyGlnTyr-H)₂ also denoted (HO-Seq. ID 1-COCH₂)₂CH-NH-Lys(Seq. ID 2-H)₂ [LPA-V]

15

FmocNCH(CH₂CO-ProValValAlaGluSerProLysLysPro)₂-chlorotrityl resin (55 mg; s = 0.44), was deprotected with 20% piperidine-DMF and coupled with Fmoc-Lys(Fmoc)-OH activated with TBTU to give the above Fmoc-Lys(Fmoc)-NHCH(CH₂CO-ProValValAlaGluSerProLysLysPro)₂-chlorotrityl resin. After deprotection of Fmoc-groups the amino acids according to the sequence (ESAT-6, 51-70) were coupled as the Fmoc protected amino acids using HOAt and DIC as coupling agents in 4 times excess with ≥2 h couplings followed by a double coupling of ≥30 min. After completed assembly of the peptide chains, the peptide-resin was washed with DMF and diethyl ether and dried in vacuo. The assembled product was cleaved from the resin with 95% TFA and 5% triisopropylsilane and precipitated by addition of ether, redissolved in acetic acid/water and lyophilised to give 2 mg of the target product C₃₀₅H₄₈₁N₈₃O₉₇ (Mw 6862.7). HPLC showed a major peak. The product was analysed for homogeneity by amino acid analysis. Alanine and leucine was used for comparison giving 3941 and 1937 pmol respectively corresponding to a Ala/Leu ratio of

20

25

30

35

2.04 (exp. 2.00). A double analysis similarly gave 2.03. The product was analysed by N-terminal amino acid sequencing and all 20 amino acids according to the ESAT sequence (Seq. ID 2) were recorded. The *Borrelia* OspC
5 sequence (Seq. ID 1) is N-terminally coupled and consequently not sequenced.

EXAMPLE 6

10 Synthesis of (HO-ProLysLysProSerGluAlaValValPro-COCH₂)₂N-Lys(AlaSerAlaAlaAlaGluIleGlyAlaPheAsnTrpGlnGlnGluThrMet-H)₂ (IDA-derivative of Seq. ID 1 extended with Seq. ID 3 on lysine)

15 Product: (HO-ProLysLysProSerGluAlaValValPro-COCH₂)₂N-Lys-(AlaSerAlaAlaAlaGluIleGlyAlaPheAsnTrpGlnGlnGluThrMet-H)₂ also denoted (HO-Seq. ID 1-COCH₂)₂N-Lys(Seq. ID 3-H)₂ [LPA-VI]

20 FmocN(CH₂CO-ProValValAlaGluSerProLysLysPro)₂-chlorotrityl resin (55 mg; s = 0.44), was deprotected with 20% piperidine-DMF and coupled with Fmoc-Lys(Fmoc)-OH activated with TBTU to give Fmoc-Lys(Fmoc)-N(CH₂CO-ProValValAlaGluSerProLysLysPro)₂-chlorotrityl resin. After
25 deprotection of Fmoc-groups with 20% piperidine-DMF the amino acids according to the ESAT-6, 1-17 sequence were coupled as the Fmoc protected amino acids using HOAt and DIC as coupling agents in 4 times excess with ≥2 h couplings followed by a double coupling of ≥30 min. After
30 completed assembly of the peptide chains, the peptide-resin was washed with DMF and diethyl ether and dried in vacuo. The product was cleaved from the resin with 95% TFA and 5% triisopropylsilane and precipitated by
35 lyophilised to give 8.7 mg of the target product. MALDI TOF MS analysis: Calculated for C₂₆₄H₄₀₉N₆₉O₈₃S₂, 5941.7;

found 5944.7 The product was analysed for homogeneity by amino acid analysis. Proline and valine from the *Borrelia* pepC10 sequence was used for comparison with Gly from the ESAT-6, 1-17 sequence. Amino acid analysis gave 5044, 2823 and 1408 pmol for Pro, Val and Gly respectively giving a Gly/Pro ratio of 0.28 (exp. 0.33) and a Gly/Val ratio of 0.50 (exp. 0.500). A double analysis similarly gave 0.27 and 0.55. The product was analysed by N-terminal amino acid sequencing and all 17 amino acids according to the ESAT-6, 1-17 sequence (Seq. ID 3) were recorded. The *Borrelia* OspC sequence (Seq. ID 1) is N-terminally coupled and consequently not sequenced.

EXAMPLE 7

15

Synthesis of $\text{CH}_2(\text{CH}_2\text{CO}-\beta\text{-Ala}-\beta\text{-AlaLysGluProAsnLysGlyValAsnProAspGluVal}\beta\text{Ala})_2$, (Glutaric acid-derivative, Fmoc- β -Ala and Dde- β -Ala synthesis of Seq. ID 4)

20 Product: $\text{CH}_2(\text{CH}_2\text{CO}-\beta\text{-Ala}-\beta\text{-AlaLysGluProAsnLysGlyValAsnProAspGluVal}\beta\text{Ala})_2$ also denoted $\text{CH}_2(\text{CH}_2\text{CO}-\beta\text{-Ala}-\beta\text{-AlaSeq. ID 4-}\beta\text{-Ala})_2$ [LPA-VII]

The $\beta\text{-Ala}-\beta\text{-AlaLysGluProAsnLysGlyValAsnProAspGluVal}\beta\text{-Ala}$ sequence (LysGluProAsnLysGlyValAsnProAspGluVal of *Chlamydia trachomatis* DnaK 357-368) was assembled on chlorotriptyl resin and on TentaGel S NH_2 resin with HMPA linker using standard coupling procedures with PyAOP 1.1 eq. and DIEA 2.2. eq. as coupling agents in DMF. In both cases the N-terminal Fmoc- α -amino group was deprotected with piperidine-DMF and subsequently coupled with a mixture of Fmoc- β -Ala and Dde- β -Ala to give an orthogonal 1:1 Fmoc/Dde-protection of the N-terminus resin-peptide chains. This was achieved by coupling with 0.75 eq. Fmoc- β -Ala and 0.25 eq. Dde- β -Ala using 1.1 eq. PyAOP and 2.2 eq. DIEA in NMP. The Fmoc-protecting group was

deprotected with 20% piperidine-DMF followed by coupling with excess glutaric acid (6 eq.) and DIEA (12 eq.) in NMP over night. The remaining N-terminal Dde- α -protecting groups were deprotected with 2% hydrazine in NMP for 10 minutes, followed by washing with 5% DIEA in NMP. Assembly was then achieved over night by activation with 6 eq. PyAOP and 12 eq. DIEA in NMP. The resulting product was cleaved from the resin with TFA-H₂O-TIS (92.5:5:2.5, vol/vol/vol). A single product with the correct mass was obtained with both resins. MALDI TOF MS: Calc. for C₁₃₅H₂₁₈N₃₈O₅₀, 3173.4; found 3169.7. (trityl-resin). Figure 9 shows the MALDI-TOF MS spectrum of the product.

EXAMPLE 8

15

Synthesis of HC(CH₂CO-LysGluProAsnLysGlyValAsnProAspGluVal β Ala)₂COOH (TCA-derivative, CH(CH₂COOH)₂COOH synthesis of Seq. ID 4)

20 Product: HC(CH₂CO-LysGluProAsnLysGlyValAsnProAspGluVal β Ala)₂COOH also denoted HC(CH₂CO-Seq. ID 4- β Ala)₂COOH [LPA-VIII]

The LysGluProAsnLysGlyValAsnProAspGluVal β Ala sequence was assembled on Tentagel S NH₂ resin with HMPA linker (s = 0.19 mmol/g; 0.02 g) using standard coupling procedures with 3 eq. DIC and 3 eq. HOAt as coupling agents in DMF. The N-terminal Fmoc- α -amino group was deprotected with piperidine-DMF and coupled with 0.55 eq. of CH(CH₂COOH)₂COOH, 1.65 eq. PyAOP and 3.3 eq. DIEA in DMF over night. The resulting product was cleaved from the resin with TFA-H₂O-TIS-EDT (90:5:2.5:2.5, vol/vol/vol/-vol) and submitted to gel filtration on G-25 Sephadex with 1% AcOH as eluent. The resulting fractions were lyophilysed to give 2 mg of the pure main product, HC(CH₂CO-LysGluProAsnLysGlyValAsnProAspGluVal β Ala)₂COOH

(approx. 20% yield). MALDI TOF MS: Calc. for $C_{124}H_{198}N_{34}O_{48}$, 2933.1; found 2932.6. Figure 10 shows the MALDI-TOF MS spectrum for the product, and Figure 11 shows the HPLC chromatogram.

5

EXAMPLE 9

Synthesis of Fmoc-NHCH(CH₂CO-AspArgValTyrIleHisProPheHisLeu-NH₂)₂, (AGA-3-derivative of angiotensin-I amide, Fmoc-3-AGA synthesis of Seq. ID 5)

10

Product: Fmoc-NHCH(CH₂CO-AspArgValTyrIleHisProPheHisLeu-NH₂)₂ also denoted Fmoc-NHCH-(CH₂CO-Seq. ID 5-NH₂)₂ [LPA-IX]

15

The peptide sequence AspArgValTyrIleHisProPheHisLeu was assembled on TentaGel S RAM resin using standard coupling procedures with DIC and HOAt as coupling agent in DMF. The N-terminal Fmoc- α -amino group was deprotected with piperidine-DMF and coupled with 0.55 eq. Fmoc-AGA, 3.3 eqv TBTU, 3.3 eqv. HOBt and 4.95 eqv. DIEA in NMP over night. The assembled product was cleaved from the resin with TFA-H₂O-triisopropylsilane (90:5:5, vol/vol/vol) to give Fmoc-NHCH(CH₂CO-AspArgValTyrIleHisProPheHisLeu-NH₂)₂ as the major product according to MS. MALDI TOF MS: Calc. for $C_{144}H_{195}N_{37}O_{30}$, 2924.4; found 2923.6.

20

25

EXAMPLE 10

Synthesis of Fmoc-NHCH(CH₂CO-AspArgValTyrIleHisProPheHisLeu-NH₂)₂ (AGA-3-derivative of angiotensin-I amide, Fmoc-3-AGA synthesis of Seq. ID 5)

30

Product: Fmoc-NHCH(CH₂CO-AspArgValTyrIleHisProPheHisLeu-NH₂)₂ also denoted Fmoc-NHCH(CH₂CO-Seq. ID 5-NH₂)₂ [LPA-IX]

35

The peptide sequence AspArgValTyrIleHisProPheHisLeu was assembled on TentaGel S RAM resin (AM-linker) using standard coupling procedures with 3 eq. DIC and 3 eq. HOAt as coupling agent in DMF. The N-terminal Fmoc- α -amino group was deprotected with piperidine-DMF and coupled with 0.55 Fmoc-AGA, 3 eqv. PyAOP, 6 eqv. DIEA in DCM/NMP (3:1 vol/vol) dissolving Fmoc-AGA in NMP, PyAOP (activating agent) in DCM and adding DIEA. Coupling over night. The assembled product was cleaved from the resin with TFA-H₂O-triisopropylsilane (90:5:5, vol/vol/vol) to give Fmoc-NHCH(CH₂CO-AspArgValTyrIleHisProPheHisLeu-NH₂)₂ as the major product according to MS. MALDI TOF MS: Calc. for C₁₄₄H₁₉₅N₃₇O₃₀, 2924.4; found 2920.8. The product may be further purified by preparative HPLC or Gel filtration if desired. Figure 12 shows the MALDI-TOF MS spectrum for the crude product. Figure 13 shows the HPLC chromatogram for the crude product.

EXAMPLE 11

Synthesis of Aloc-NHCH(CH₂CO-AspArgValTyrIleHisProPheHisLeu-NH₂)₂ (AGA-3-derivative of angiotensin-I amide, Fmoc- β -Ala and Dde- β -Ala synthesis of Seq. ID 5 using orthogonal protection of the sequence to be assembled)

Product: Aloc-NHCH(CH₂CO-AspArgValTyrIleHisProPheHisLeu-NH₂)₂ also denoted Aloc-NHCH-(CH₂CO-Seq. ID 5-NH₂)₂ [LPA-X]

30

The peptide sequence AspArgValTyrIleHisProPheHisLeu was assembled on TentaGel S RAM resin using standard coupling procedures with 3 eq DIC and 3 eq. HOAt as coupling agents in DMF. The N-terminal Fmoc- α -amino group was deprotected with piperidine-DMF and subsequently coupled with a mixture of Fmoc- β -Ala and Dde- β -Ala to give an

35

orthogonal 1:1 Fmoc/Dde-protection of the N-terminus resin-peptide chains. The Fmoc-protecting group was deprotected with 20% piperidine-DMF followed by over night coupling with excess Aloc-3-AGA (6 eq.) and DIEA (12 eq.) in NMP. The remaining N-terminal Dde- α -amino groups were deprotected with 2% hydrazine in NMP for 10 minutes, followed by washing with 5% DIEA in NMP. Assembly was then achieved over night by activation with 6 eq. PyAOP and 12 eq. DIEA in NMP. The resulting product was cleaved from the resin with TFA-H₂O-TIS (92.5:5:2.5, vol/vol/vol). MALDI TOF MS: Calc. for C₁₃₉H₁₉₉N₃₉O₃₂, 2928.6; found 2927.5.

EXAMPLE 12

15

Synthesis of Fmoc-AspProThrGlnAsnIleProProGly-NHCH(CH₂CO-AspArgValTyrIleHisProPheHisLeu-NH₂)₂ (AGA-3-derivative of angiotensin-I amide extended with *Chlostridium thermosaccharolyticum* peptide sequence 19-27)

20

Product: Fmoc-AspProThrGlnAsnIleProProGly-NHCH(CH₂CO-AspArgValTyrIleHisProPheHisLeu-NH₂)₂ also denoted Fmoc-Seq. ID 6-NHCH(CH₂CO-Seq. ID 5-NH₂)₂ [LPA-XI]

25

The *Chlostridium thermosaccharolyticum* peptide sequence 19-27 AspProThrGlnAsnIleProProGly was synthesised on chlorotrityl resin using standard peptide synthesis and cleaved from the resin with TFA-DCM (1:99, vol/vol) affording the fragment Fmoc-Asp(tBu)ProThr(tBu)Gln(Trt)-Asn(Trt)IleProProGly-OH. FmocNHCH(CH₂CO-AspArgValTyrIleHisProPheHisLeu-NH₂)₂ was assembled as above (Example 11) and coupled with 3 eq. of the fragment using 3 eq. PyAOP and 6 eq. DIEA as coupling agents in NMP over night. The resulting fragment coupled product was cleaved from the resin with TFA-H₂O-TIS (90:5:5, vol/vol/vol) to give the target compound together with some NH₂CH(CH₂CO-AspArgVal-

30

35

TyrIleHisProPheHisLeu-NH₂)₂. MALDI TOF MS: Calc. for C₁₈₄H₂₅₄N₄₆O₄₆, 3846.3; found 3842.8.

EXAMPLE 13

5

The performance of an ELISA based on LPA-I.

The capability of LPA-I according to the present invention to react with sera from patients with neuroborreliosis was evaluated against that of an ELISA
10 based on biotinylated-pepC10, wherein pepC10 is identical to the amino acid sequence of Seq. ID 1.

Indirect IgM-ELISA using LPA-I and biotinylated-pepC10

Preparing plates for the LPA-I-ELISA: Microtiter plates
15 (Maxisorb; Nunc, Roskilde, Denmark) were coated overnight at 4°C with 0.6 µg/ml of LPA-I diluted in 0.05 M bicarbonate pH 9.6, blocked with 3% (w/v) milk powder in phosphate-buffered saline (PBS) for 1 h, and reacted with sera diluted 1/200 in 1% (w/v) milk powder in PBS
20 containing 0.05% Tween 20 (PBST) and 0.7 M NaCl for 2 h. The secondary antibody was a peroxidase conjugated rabbit anti-human IgM (Dako, Denmark, codes P-215) diluted 1/1000 in 1% (w/v) milk powder in PBST. After 1 h of incubation, bound secondary antibody was quantitated by
25 colouring with o-phenylenediamine and H₂O₂ in citrate buffer (Sigma, St. Louis, Mo.) for 30 min. Optical density (OD) at 492 nm was determined in a plate reader (EAR 400 AC-SLT, Labinstruments, Austria). Plates were washed extensively with PBST-0.5M NaCl between each
30 incubation step. The 98% specific diagnostic cut-off level was OD of 0.270 for IgM detection.

Preparing plates for the biotinylated-pepC10-ELISA:
ELISAs with biotinylated-pepC10, were performed
35 essentially as described above for LPA-I, except that microtiter plates were first coated with streptavidin

(2.5 µg/ml) (Zymed, S. Avidinii) in citrate buffer (pH 5) overnight at 4°C, and then incubated with biotinylated pepC10 (0.1 µg/ml) in 1% (w/v) milk powder in PBST-0.37 M NaCl overnight at 4°C.

5

Serum specimens from patients

(i) Sera from 97 Danish patients with neuroborreliosis (NB).

All patients had been hospitalised in 1994 (58 males and
10 42 females between 4 and 80 years of age; median age, 49). The selection was based on a positive test for *B. burgdorferi* specific intrathecal antibody synthesis. The diagnosis was further corroborated by the presence of lymphocytic pleocytosis in CSF (median cell count,
15 124/µl) and one of the following manifestations: lymphocytic meningoradiculitis with typical painful radiculitis, mononeuritis multiplex (Bannwarths syndrome), radiculomyelitis, monosymptomatic fascialis paresis (all children), subacute lymphocytic meningitis, and chronic
20 progressive encephalomyelitis. Of the 100 patients, 48 recalled a previous EM-like skin lesion. Disease duration, defined as the time from onset of neurological symptoms until diagnostic blood and CSF samples were taken, ranged from 1 week to 26 weeks (median duration, 3
25 weeks).

Control serum specimens

Sera from 80 healthy controls were used for determination of the 98%-specific cut-off level in ELISAs.

30

Results

Comparing results of the LPA-I ELISA with results of the biotinylated-pepC10 ELISA.

35 The following table (Table 1) shows the frequency (%) of positive Lyme borreliosis sera in the early stages of

Lyme borreliosis found by the above-described LPA-I and biotinylated-pepC10 ELISAs.

TABLE 1. IgM reactivity of NB sera to biotinylated-pepC10 and LPA-I.

	No. (%) of IgM reactive sera to	
	biotinylated pepC10	LPA-I
NB (n=97)	n=42 (43.3%)	n=41 (42.3%)

It can be concluded that the two assays have nearly the same sensitivity.

Figure 7 compares the individual results obtained in patients with NB regarding the quantitative measurement of IgM in the two ELISAs. The vertical and horizontal lines marks the 98% specific diagnostic cut-off levels for the respective LPA-I and biotinylated-pepC10 ELISA (0.275 and 0.460). As can be seen the OD titres are of the same magnitude in both assays.

It can be concluded that in this set-up, the sensitivity of the two assays are comparable. Further since the OD cut-off-value in the pepC10 ELISA is higher than the OD cut-off-value in the LPA-I ELISA it is expected that the LPA-I ELISA will perform better than the biotinylated-pepC10-ELISA when testing a large number of serum samples. Importantly, three serum samples in Figure 7 tested positive in the biotinylated-pepC10 ELISA but negative in the LPA-I ELISA. The positive reaction of all three sera is not due to anti-Borrelia antibodies but is due to the presence of anti-streptavidin antibodies which reacted with streptavidin in the biotinylated-pepC10 ELISA only. The present invention, therefore, is superior

to ELISA procedures which employ peptides coupled to streptavidin because it reduces the number of false-positive serum samples.

5 EXAMPLE 14

Immunogenicity of the LPA-VI construct.

The capability of an LPA according to the present invention to induce a humoral immune response was studied
10 by immunising mice with LPA-VI. Five female C57 black mice (age 6 to 8 weeks) were immunised two times intraperitoneally with a 2-week interval and were bled on days 0, 14, and 28. Each immunisation dose contained 5 µg of LPA-VI dissolved in 0.25 ml 0.9% NaCl and supplemented
15 with 1 mg Al(OH)₃ (Superfos Biosector, Denmark) and 0.25 ml Freund's incomplete adjuvant. The serum samples were diluted 200 fold and then tested by ELISA for Ig reactivity against OspC produced on recombinant form (WO 97/42221) and LPA-I (Figure 8). It is concluded that LPA-
20 VI is highly effective in inducing antibodies against the C-terminal B-cell epitope of OspC since the mice generated antibodies which recognise both recombinant OspC and LPA-I.

25 Figure 8. Mean ELISA titer of sera from five mice immunised with LPA-VI on days 0 and 14. The microdilution plates were coated with LPA-I (diamond) or recombinant OspC (Squares), and the sera were analysed in a 200-fold dilution.

SEQUENCE LISTING

<110> HOLM, Arne and Statens Serum Institut

5 <120> Method for preparing and Ligand Presenting Assembly
(LPA), and LPA, and uses thereof

<130> P199801104 WO

10 <140>

<141>

<150> DK PA 1998 01233

<151> 1998-09-29

15

<160> 6

<170> PatentIn Ver. 2.1

20 <210> 1

<211> 10

<212> PRT

<213> Artificial Sequence

25 <220>

<223> Description of Artificial Sequence: Sequence
derived from the OspC protein of Borrelia
burgdorferi

30 <400> 1

Pro Val Val Ala Glu Ser Pro Lys Lys Pro

1

5

10

35 <210> 2

<211> 20

<212> PRT

<213> Artificial Sequence

<220>

5 <223> Description of Artificial Sequence: ESAT-6, 51-70
sequence of Mycobacterium tuberculosis

<400> 2

Gln Leu Ala Asn Asn Leu Glu Thr Ala Thr Ala Asp Trp Lys Gln Gln
10 1 5 10 15

Val Gly Gln Tyr
20

15

<210> 3

<211> 17

<212> PRT

<213> Artificial Sequence

20

<220>

<223> Description of Artificial Sequence: ESAT-6, 1-17
sequence of Mycobacterium tuberculosis

25 <400> 3

Ala Ser Ala Ala Ala Glu Ile Gly Ala Phe Asn Trp Gln Gln Glu Thr
1 5 10 15

Met

30

<210> 4

<211> 12

35 <212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Chlamydia
trachomatis DnaK 357-368 sequence

5

<400> 4

Lys Glu Pro Asn Lys Gly Val Asn Pro Asp Glu Val

1 5 10

10

<210> 5

<211> 10

<212> PRT

<213> Artificial Sequence

15

<220>

<223> Description of Artificial Sequence: Angiotensin I
sequence

20

<400> 5

Asp Arg Val Tyr Ile His Pro Phe His Leu

1 5 10

25

<210> 6

<211> 9

<212> PRT

<213> Artificial Sequence

30

<220>

<223> Description of Artificial Sequence: Clostridium
thermosaccharolyticum peptide sequence 19-27

<400> 6

35 Asp Pro Thr Gln Asn Ile Pro Pro Gly

1

5

REFERENCES

1. PCT/US90/02039 (Tam)
- 5 2. Tam, J.P. (1996) J. Immun. Methods 196, 17-32.
3. Tomalia, D.A., Naylor, A.M., and Goddard III, W.A. (1990) Angew. Chem. Int. Ed. Engl. 29, 138-175.
- 10 4. Hodge, P. (1993) Nature (London) 362, 18-19.
5. Tam, J.P. (1995) Chapter 11, Epitope Mapping with Peptides in Peptides: Synthesis, Structures, and Applications (B. Gutte, ed.), Academic Press, 1995.
- 15 6. Heegaard, P.M.H., Bøg, Y.S., Jakobsen, P.H., and Holm, A. (1997) In press.
7. Sarin et al., Anal. Biochem. 117, 147-157 (1981)
- 20 8. Nardelli et al., J. Immunol. 148, 914-920 (1992b)
9. Nardelli et al., Vaccine 12, 1335-1339 (1994)
- 25 10. Keah, H.H., Kecorius, E., and Hearn, M.T.W. (1998) J. Peptide Res. 51, 2-8.
11. Lu, Y.-A., Clavijo, P., Galantino, M., Shen, Z.-Y. and Tam, J.P. (1991) Mol. Immunol. 28, 623-630.
- 30 12. Defoort, J.P., Nardelli, B., Huang, W. and Tam, J.P. (1992) Int. J. Pept. Prot. Res. 40, 214-221.
13. Drijfhout, J. W. and Bloemhoff, W. (1991) Int. J. Pept. Prot. Res. 37, 27-32.
- 35

14. WO 93/24523 (Hafslund Nycomed)

15. Lange, M., Cuthbertson, A.S., Towart, R. and Fischer, P.M. (1998) J. Peptide Science 4, 289-293.

5

16. Bhatnagar, P.K., Agner, E.K., Alberts, D., Arbo, B.E., Callahan, J.F., Cuthbertson, A.S., Engelsen, S.J., Fjordingstad, H., Hartmann, M., Heerding, D., Hiebl, J., Huffman, W.F., Hysben, M., King, A.G., Kremminger, P.,
10 Kwon, C., LoCastro, S., Løvhaug, D., Pelus, L.M., Petteway, S. and Takata, J.S., (1996) J. Med. Chem. 39, 3814-3819.

17. Alberts et al., Peptides: Chemistry, Structure and
15 Biology (Hodges, R.S., and Smith, J.A., Eds.), 367-369 (1993), ESCOM, Leiden

18. Tuchscherer, G., and Mutter, M., Journal of Biotechnology 41, 197-210 (1995)

20

19. WO 94/02506 (The School of Pharmacy, University of London)

20. WO 98/32469 (Nycomed Imaging AS)

25

21. Marshall, G. R., Tetrahedron 49, 3547-3558 (1993)

22. WO 98/11125 (Arne Holm et al.)

30

23. WO 98/15648

24. Peptide Synthesis Protocols, Methods in Molecular Biology Vol. 35, Chapter 15, 303-316 in Peptide Synthesis via Fragment Condensation by R. Nyfeler, M.W. Pennington
35 and B.M. Dunne Eds., Humana Press, 1994

25. WO 97/42221 (Statens Seruminstitut)

26. WO 95/01441 (Statens Seruminstitut)

5 27. US 4 608 251

28. US 4 601 903

29. US 4 599 231

10

30. US 4 599 230

31. US 4 596 792

15 32. US 4 578 770

CLAIMS

1. A method for preparing a ligand presenting assembly
(LPA) enabling presentation of desired sequence(s)
5 comprising the steps of
- (a) providing by solid phase synthesis or fragment
coupling ligands comprising desired sequence(s),
the ligands being attached to a solid phase,
10
- (b) if necessary, deprotecting any N-terminal amino
groups while the ligand(s) are still attached to
the solid phase,
- 15 (c) reacting the ligand(s) having unprotected N-
terminal amino groups with an achiral di-, tri- or
tetracarboxylic acid so as to provide a construct
having a ring structure, and
- 20 (d) cleaving the construct from the solid phase so as
to provide an LPA comprising ligands having free C-
terminal groups.
2. A method according to claim 1 for the preparation of
25 an LPA enabling presentation of desired sequence(s) or
desired sequence(s) and chemical moieties further
comprising the steps of
- (c¹) if present, prior to step (d), deprotecting any N-
30 protected amino groups originating from the
carboxylic acid used in step (c),
- (c²) continuing the solid phase synthesis or fragment
coupling so as to provide ligand(s) comprising
35 desired sequence(s) having at least one N-protected

N-terminal amino group and/or attaching chemical moieties, and

5 (c³) if present, deprotecting any N-terminal amino group(s) prior to or after step (d).

3. The method according to claim 1 or 2, wherein the achiral acid used in step (c) is of the general formula

10 $X[(A)_n\text{COOH}][(B)_m\text{COOH}]$

wherein n and m independently are an integer of from 1 to 20, X is HN, A and B independently are a substituted or unsubstituted C₁₋₁₀ alkyl, a substituted or unsubstituted C₂₋₁₀ alkenyl, a substituted or unsubstituted cyclic moiety, a substituted or unsubstituted heterocyclic moiety, or a substituted or unsubstituted aromatic moiety, or A and B together form a substituted or unsubstituted cyclic moiety, a substituted or unsubstituted heterocyclic moiety, or a substituted or unsubstituted aromatic moiety, or

n and m are 0 or an integer of from 1 to 20, X is H₂N(CR₂)_pCR, or RHN(CR₂)_pCR, wherein p is 0 or an integer of from 1 to 20, wherein each R is H, a substituted or unsubstituted C₁₋₁₀ alkyl, a substituted or unsubstituted C₂₋₁₀ alkenyl, a substituted or unsubstituted cyclic moiety, a substituted or unsubstituted heterocyclic moiety, or a substituted or unsubstituted aromatic moiety, and A and B are both a substituted or unsubstituted C₁₋₁₀ alkyl, a substituted or unsubstituted C₂₋₁₀ alkenyl, a substituted or unsubstituted cyclic moiety, a substituted or unsubstituted heterocyclic moiety, or A and B together form a substituted or unsubstituted cyclic moiety, a substituted or

unsubstituted heterocyclic moiety, or a substituted or unsubstituted aromatic moiety, or

n and m are 0 or an integer of from 1 to 20, X is
5 HO(CR₂)_pCR, HS(CR₂)_pCR, halogen-(CR₂)_pCR, HOOC(CR₂)_pCR,
ROOC(CR₂)_pCR, HCO(CR₂)_pCR, RCO(CR₂)_pCR, or [HOOC(A)_n]-
[HOOC(B)_m]CR(CR₂)_pCR, wherein p is 0 or an integer of
from 1 to 20, each R independently is H, a substituted or
unsubstituted C₁₋₁₀ alkyl, a substituted or unsubstituted
10 C₂₋₁₀ alkenyl, a substituted or unsubstituted cyclic
moiety, a substituted or unsubstituted heterocyclic
moiety, or a substituted or unsubstituted aromatic
moiety, and A and B are both a substituted or
unsubstituted C₁₋₁₀ alkyl, a substituted or unsubstituted
15 C₂₋₁₀ alkenyl, a substituted or unsubstituted cyclic
moiety, a substituted or unsubstituted heterocyclic
moiety, or A and B together form a substituted or
unsubstituted cyclic moiety, a substituted or
unsubstituted heterocyclic moiety, or a substituted or
20 unsubstituted aromatic moiety, or

n and m are 0 or an integer of from 1 to 20, X is
H₂N(CR₂)_p, RHN(CR₂)_p, HO(CR₂)_p, HS(CR₂)_p, halogen-(CR₂)_p,
HOOC(CR₂)_p, ROOC(CR₂)_p, HCO(CR₂)_p, RCO(CR₂)_p or
25 [HOOC(A)_n][HOOC(B)_m], wherein p is 0 or an integer of
from 1 to 20, each R independently is H, a substituted or
unsubstituted C₁₋₁₀ alkyl, a substituted or unsubstituted
C₂₋₁₀ alkenyl, a substituted or unsubstituted cyclic
moiety, a substituted or unsubstituted heterocyclic
30 moiety, or a substituted or unsubstituted aromatic
moiety, and A and B together form a substituted or
unsubstituted cyclic moiety, a substituted or
unsubstituted heterocyclic moiety, or a substituted or
unsubstituted aromatic moiety.

4. The method according to any one of claims 1-3, wherein the achiral acid is a di- or tricarboxylic acid.

5. The method according to any one of claims 1-4, wherein
5 the achiral acid is selected among imino diacetic acid, 2-amino malonic acid, 3-amino glutaric acid, 3-methylamino glutaric acid, 3-chloro glutaric acid, 3-carboxymethyl glutaric acid, 3-methoxy-carbonyl glutaric acid, 3-acetyl glutaric acid, glutaric acid,
10 tricarballic acid, 3,4-bis-carboxymethyl adipic acid, 4-(2-carboxyethyl)-pimelic acid, (3,5-bis-carboxymethyl-phenyl)-acetic acid, 3,4-bis-carboxymethyl-adipic acid, benzene-1,2,4,5-tetra carboxylic acid, 4-(3-carboxy-allylamino)-but-2-enoic acid, 4,4-imino-dibenzoic acid,
15 1,4-dihydropyridine-3,5-dicarboxylic acid, 5-amino isophthalic acid, 2-chloro malonic acid, 3-hydroxy glutaric acid, and benzene-1,3,5-tricarboxylic acid.

6. The method according to any one of claims 1-5, wherein
20 the desired sequence is a peptide comprising naturally occurring amino acids and/or non-naturally occurring amino acids, a peptide nucleic acid (PNA) sequence or a peptidomimetic.

25 7. The method according to any one of claims 2-6, wherein the chemical moiety is an entity enhancing the solubility or immunogenicity of the LPA obtained according to claims 1-5, or is an entity suitable for directing the LPA to its target, or a marker.

30

8. The method according to claim 7, wherein the chemical moiety is selected from fatty acids, antibodies or peptides for directing the LPA to its target, fluorophores, biotin, enzymes such as horse radish
35 peroxidase, alkaline phosphatase and soya bean peroxidase, or nucleic acid sequences.

9. The method according to any one of claims 1-8, wherein the desired sequence comprises all or part of one or more B cell epitopes, all or part of one or more T cell epitopes, or all or part of one or more B and T cell epitopes, or mimics thereof.

10. The method according to claim 9, wherein at least one of the sequences is derived from a sequence, wherein the C-terminal amino acids are important for an immune response.

11. The method according to any one of claims 1-10, wherein the desired sequence is derived from viruses, bacteria, toxins, allergens, autoimmune system-related compounds, cancer related compounds, cell adhesion molecules, neurotropic factors, fungi or parasites, or a sequence homologous thereto.

12. The method according to any one of claims 1-11, wherein the desired sequence is derived from OspC protein of *Borrelia burgdorferi* or is a homologous sequence capable of reacting with anti-OspC antibodies or of provoking an immune response.

13. The method according to claim 12, wherein the LPA obtained provide a C-terminal presentation of the C-terminal sequence Pro-Lys-Lys-Pro of OspC.

14. The method according to any one of claims 1-11, wherein the desired sequence is derived from the flagellum of *Borrelia burgdorferi* or is a homologous sequence capable of reacting with anti-flagellum antibodies.

15. The method according to claim 12 or 13, wherein the LPA obtained provide C-terminal presentation of sequence(s) derived from OspC of *Borrelia burgdorferi* and further comprises desired sequence(s) derived from the flagellum of *Borrelia burgdorferi*.

16. The method according to any one of claims 1-11, wherein the desired sequence is derived from *Mycobacterium tuberculosis*.

17. The method according to claim 12 and 13, wherein the LPA obtained further comprises desired sequence(s) derived from *Mycobacterium tuberculosis*.

18. The method according to claim 16 or 17, wherein the desired sequence comprises the ESAT-6, 51-70 sequence protein or the ESAT-6, 1-17 sequence protein of *Mycobacterium tuberculosis*.

19. An LPA obtainable by the method defined in claims 1-18.

20. An LPA according to claim 19, wherein the ligands comprising the desired sequence is selected from a peptide comprising naturally occurring and/or non-naturally occurring amino acids, a PNA sequence or a peptidomimetic.

21. An LPA according to claim 19 or 20, wherein the chemical moiety is an entity enhancing the solubility or immunogenicity of the LPA of claims 1-20 or is an entity suitable for directing the LPA to its target, or a marker.

22. An LPA according to any one of claims 19-21, wherein the chemical moiety is selected from fatty acids,

antibodies or peptides for directing the LPA to its target, fluorophores, biotin, enzymes such as horse radish peroxidase, alkaline phosphatase and soya bean peroxidase, or nucleic acid sequences.

5

23. An LPA according to any one of claims 19-22, wherein the desired sequence comprises all or part of one or more B cell epitopes, all or part of one or more T cell epitopes, or all or part of one or more B and T cell epitopes, or mimics thereof.

10

24. An LPA according to claim 23, wherein at least one of the sequences is derived from a sequence, wherein the C-terminal amino acids are important for an immune response.

15

25. An LPA according to any one of claims 19-24, wherein the desired sequence is derived from viruses, bacteria, toxins, allergens, auto-immune system-related compounds, cancer related compounds, fungi or parasites, or a sequence homologous thereto.

20

26. An LPA according to any one of claims 19-24, wherein the desired sequence is derived from the OspC protein of *Borrelia burgdorferi*, or is a homologous sequence capable of reacting with anti-OspC antibodies or of provoking an immune response.

25

27. An LPA according to claim 26, wherein the LPA obtained provide a C-terminal presentation of the C-terminal sequence Pro-Lys-Lys-Pro of OspC.

30

28. An LPA according to any one of claims 19-27 selected from

35

[LPA-I]: FmocN(CH₂CO-ProValValAlaGluSerProLysLysPro-OH)₂,

[LPA-II]: biotin-NH(CH₂)₅CON(CH₂CO-ProValValAlaGluSerPro-LysLysPro-OH)₂,

5 [LPA-III]: NH₂CH(CH₂CO-ProValValAlaGluSerProLysLysPro-OH)₂, and

[LPA-IV]: H-Lys-NHCH(CH₂CO-ProValValAlaGluSerProLysLys-Pro-OH)₂.

10

29. An LPA according to any one of claims 19-25, wherein the desired sequence is derived from the flagellum of *Borrelia burgdorferi* or is a homologous sequence capable of reacting with anti-flagellum antibodies.

15

30. An LPA according to any one of claims 26-28 further comprising desired sequence(s) derived from the flagellum of *Borrelia burgdorferi* or is a homologous sequence capable of reacting with anti-flagellum antibodies.

20

31. An LPA according to any one of claims 19-25, wherein the desired sequence is derived from *Mycobacterium tuberculosis*.

25 32. An LPA according to any one of claims 26-28 further comprising desired sequence(s) derived from *Mycobacterium tuberculosis*.

30 33. An LPA according to claim 31 or 32, wherein the desired sequence comprises the ESAT-6, 51-70 sequence protein or the ESAT-6, 1-17 sequence protein of *Mycobacterium tuberculosis*.

35 34. An LPA according to any one of claims 31-33 selected from

[LPA-V]: (HO-ProLysLysProSerGluAlaValValPro-COCH₂)₂CH-NH-Lys(GlnLeuAlaAsnAsnLeuGluThrAlaThrAlaAspTrpLysGlnGln-ValGlyGlnTyr-H)₂, and

5 [LPA-VI]: (HO-ProLysLysProSerGluAlaValValPro-COCH₂)₂N-Lys(AlaSerAlaAlaAlaGluIleGlyAlaPheAsnTrpGlnGlnGluThrMet-H)₂.

10 35. An immunological composition for raising an immune response in an animal, including a human being, comprising an LPA as defined in any of claim 1-34.

36. An immunological composition according to claim 35 comprising an LPA as defined in any of claims 1-34 in
15 combination with an adjuvant.

37. An immunological composition according to claim 35 or 36 which is in the form of a vaccine.

20 38. A vaccine according to claim 37 for oral or parenteral, e.g. subcutaneous, intramuscular, intradermal, nasal, or pulmonary administration.

39. A method for generating antibodies in an animal,
25 including a human being, which method comprises administering to said mammal an antibody-generating amount of the LPA as defined in any of claims 1-34.

40. A kit for use in the diagnosis of infections caused
30 by viruses, bacteria, toxins, allergens, autoimmune system-related compounds, cancer related compounds, cell adhesion molecules, neurotropic factors, fungi or parasites, which kit comprises an LPA as defined in any of claims 1-34 together with means for detecting or
35 visualising binding between the LPA and the substance to be detected.

41. A kit for use in diagnosing diseases caused by *Borrelia burgdorferi sensu lato*, which kit comprises an LPA as defined in any of claims 26-30 or 32-34 together
5 with means for detecting or visualising binding between the LPA and the substance to be detected.

42. Use of an LPA according to any one of claims 19-34 for preparing a pharmaceutical composition for the
10 treatment, alleviation or prophylaxis of diseases caused by viruses, bacteria, toxins, allergens, autoimmune system-related compounds, cancer related compounds, cell adhesion molecules, neurotropic factors, fungi or parasites.

FIG. 1




FIG. 2




FIG. 3




FIG. 4




FIG. 5





FIG. 6






FIG. 7





FIG. 8

9/13

FIG. 9



10/13

FIG. 10




FIG. 11






FIG. 12



13/13

FIG. 13



INTERNATIONAL SEARCH REPORT

Inter nal Application No
PCT/DK 99/00510

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07K1/04 C07K7/02 C07K14/00 A61K39/04 A61K39/02
A61K39/385

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07K A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	TAM J P: "Recent advances in multiple antigen peptides" JOURNAL OF IMMUNOLOGICAL METHODS, vol. 196, no. 1, 1 January 1996 (1996-01-01), page 17-32 XP004021277 cited in the application the whole document ---	19-25, 35-40,42
X	WO 94 02506 A (GIBBONS WILLIAM ANTHONY ;TOTH ISTVAN (GB); UNIV LONDON PHARMACY (G) 3 February 1994 (1994-02-03) abstract --- -/-	19-25, 35-40,42

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "&" document member of the same patent family

Date of the actual completion of the international search

11 January 2000

Date of mailing of the international search report

18/01/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Cervigni, S

INTERNATIONAL SEARCH REPORT

Inter nal Application No
PCT/DK 99/00510

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>PELUSO S ET AL: "Protein Mimetics (TASP) by Sequential Condensation of Peptide Loops to an Immobilised Topological Template"</p> <p>TETRAHEDRON,</p> <p>vol. 53, no. 21, 26 May 1997 (1997-05-26), page 7231-7236 XP004105702</p> <p style="text-align: center;">---</p>	
X	<p>G. TUCHSCHERER ET AL: "Templates in protein de novo design"</p> <p>J. BIOTECHNOLOGY,</p> <p>vol. 41, 1995, pages 197-210, XP002104629 page 207, column 1; figure 9</p> <p style="text-align: center;">---</p>	19-25, 35-40,42
X	<p>WO 98 32469 A (GOLDING LOUISE ;NYCOMED IMAGING AS (NO); WOLFE HENRY (US); KELLAR) 30 July 1998 (1998-07-30) abstract</p> <p style="text-align: center;">---</p>	19-25,35
A	<p>M. HARBOE ET AL.: "B-cell epitopes and Quantification of ESAT-6 protein of Mycobacterium tuberculosis"</p> <p>INFECTION AND IMMUNITY,</p> <p>vol. 66, no. 2, February 1998 (1998-02), pages 717-723, XP002104630</p> <p style="text-align: center;">---</p>	
A	<p>WO 97 42221 A (STATENS SERUMINSTITUT ;MATHIESEN MARIANNE JARTVED (DK); THEISEN MI) 13 November 1997 (1997-11-13)</p> <p style="text-align: center;">-----</p>	

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/DK 99/00510

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9402506 A	03-02-1994	AU 4715493 A DE 69315842 D DE 69315842 T EP 0652896 A ES 2111167 T US 5882645 A	14-02-1994 29-01-1998 09-04-1998 17-05-1995 01-03-1998 16-03-1999
WO 9832469 A	30-07-1998	AU 5871798 A NO 993662 A	18-08-1998 28-09-1999
WO 9742221 A	13-11-1997	CA 2253374 A EP 0896583 A	13-11-1997 17-02-1999